

# Whitebark Pine Pilot Fieldwork Report

## Klamath National Forest



By Michael Kauffmann<sup>1</sup>, Sara Taylor<sup>2</sup>, Kendra Sikes<sup>3</sup>, and Julie Evens<sup>4</sup>

In collaboration with:

Marla Knight, Forest Botanist, Klamath National Forest

Diane Ikeda, Regional Botanist, Pacific Southwest Region, USDA Forest Service

January, 2014

**November 2018 Update: Page 47**

- 
1. Kauffmann, Michael E., Humboldt State University, Redwood Science Project, 1 Harpst Street, Arcata, CA 95521, [michaielekauffmann@gmail.com](mailto:michaielekauffmann@gmail.com)
  2. Taylor, Sara M., California Native Plant Society, 2707 K Street, Suite 1, Sacramento, CA 95816, [staylor@cnps.org](mailto:staylor@cnps.org)
  3. Sikes, Kendra., California Native Plant Society, 2707 K Street, Suite 1, Sacramento, CA 95816, [ksikes@cnps.org](mailto:ksikes@cnps.org)
  4. Evens, Julie., California Native Plant Society, 2707 K Street, Suite 1, Sacramento, CA 95816, [jevans@cnps.org](mailto:jevans@cnps.org)



CALIFORNIA  
NATIVE PLANT SOCIETY



Photo on cover page: *Pinus albicaulis* seen from Boulder Peak in the Marble Mountain Wilderness area, Klamath National Forest

All photos by Michael Kauffmann unless otherwise noted  
All figures by Kendra Sikes unless otherwise noted

Acknowledgements: We would like to acknowledge Marla Knight, Julie K. Nelson and Diane Ikeda for reviewing and providing feedback on this report. We also would like to thank Matt Bokach, Becky Estes, Jonathan Nesmith, Nathan Stephenson, Pete Figura, Cynthia Snyder, Danny Cluck, Marc Meyer, Silvia Haultain, Deems Burton and Peggy Moore for providing field data points or mapped whitebark pine for this project.

Special thanks to Dr. Jeffrey Kane and Jay Smith for adventuring into the northern California wilds and helping with field work.

Suggested report citation:

Kauffmann, M., S. Taylor, K. Sikes, and J. Evens. 2014. Klamath National Forest: Whitebark Pine Pilot Fieldwork Report. Unpublished report. California Native Plant Society Vegetation Program, Sacramento, CA.

<http://pacslope-conifers.com/conifers/pine/wbp/CNPS-Reports/index.html>

# Table of Contents

i	Table of Contents
ii	Figures
ii	Tables
1	Background
3	Introduction
3	Methods and Materials
4	Results
5	Forest-Specific Conclusions, Discussion, and Recommendations
7	Summary Tables from the CNDDDB Rare Plant Occurrence Forms
10	Overview and Regional Detail Maps of 2013 Locations Visited on the National Forest, including negative reports
20	Photos from 2013 Field Work
26	Literature Cited
28	Appendix 1: Inventory and Monitoring Protocols and Field Forms from 2013 including the CNPS Vegetation Rapid Assessment/Relevé Form
41	Appendix 2: Recommended Protocols for Future Work

## Figures

- 2 Figure 1: Whitebark pine range in western North America (by Michael Kauffmann)
- 10 Figure 2: Overview Maps of 2013 Locations Visited on the National Forest
- 11 Figure 3: Goosenest Ranger District - Detailed Maps
- 12 Figure 4: Goosenest Ranger District - North
- 13 Figure 5: Goosenest Ranger District - South
- 14 Figure 6: Scott River Ranger District - Overview
- 15 Figure 7: Scott River Ranger District - China Mountain Region
- 16 Figure 8: Scott River Ranger District - Marble Mountain Overview
- 17 Figure 9: Scott River Ranger District - Marble Mountain Overview
- 18 Figure 10: Scott River Ranger District - Russian Wilderness Overview
- 19 Figures 11: Range of whitebark pine in the divide between the Big Bar, Weaverville, and Salmon River Ranger Districts - Trinity Alps Wilderness
- 20 Figures 12-14: The Whaleback
- 21 Figures 18: Ash Creek Butte
- 22 Figures 19: Antelope Creek RNA
- 23 Figures 20-23: Boulder Peak region
- 24 Figures 24-25: South China Mountain

## Tables

- 7 Table 1: Whitebark pine population area by forest in northern California
- 8 Table 2: Rapid assessment summary for Klamath National Forest
- 9 Tables 3: Whitebark pine attributes from Rapid Assessments

# Background

Whitebark pine (*Pinus albicaulis*) is a long-lived and slow-growing tree found in upper montane to subalpine forests of southwestern Canada and the western United States. It regularly defines upper treeline and co-occurs with other conifers. Of the approximately 250,000 acres where whitebark pine forms pure stands in California, >95% is on public land, often in remote wilderness settings on National Forest and Park lands; however, the acreage of the pine's presence in mixed-stands across the state is much greater (see Figure 1).

Across the state, the species is found from 1,830 – 4,240 m (6,000'–13,899') in the Sierra Nevada, Cascade, Warner, and Klamath mountains where it is an outlier of a much broader range (Arno et al. 1989, Murray 2005) from the more contiguous Rocky Mountains and Cascades in western North America. Within this range, the species prefers cold, windy, snowy, and generally moist zones. In the moist areas of the Klamath and Cascades, it is most abundant on the warmer and drier sites. In the more arid Warner Mountains and in the Sierra Nevada, the species prefers the cooler north-face slopes and more mesic regions. But some of these phytogeographic patterns are shifting.

Western coniferous forests are currently undergoing large-scale changes in composition and distribution. These changes are due to shifts in the following: climate regimes, insect and fungal pathogen distributions, fire return intervals, fire severity/intensity, and logging practices—among others. High elevation five-needle pines have been harbingers for climate change for millions of years, and because high-elevation ecosystems are likely to be the first to register the impacts of global climate change (Bunn et al. 2005), surveying high elevation five-needle pine is a way to catalog trends in vegetation and climatic shifts.

Unlike other five-needle pines, whitebark pine is set apart in that its cone does not open at maturity and its seed is “wingless”; consequently, they are solely dependent on Clark's nutcrackers (*Nucifraga columbiana*) for seed distribution and future seedling recruitment. The birds open the cone, collect the seeds, and cache them. Inevitably, around 20% of the seeds are forgotten or moved by other animals (Lanner 1996) and, in the years following, clumps of whitebark pine saplings grow from these forgotten caches. These two species are keystone mutualists, where the loss of one species would have a profound impact upon the ecosystem as a whole.

Whitebark pine (WBP) is currently the most susceptible of the five-needle pines to mortality due to the combined effects of climate change-induced disturbance. Mortality across much of its range is attributed to white pine blister rust (WPBR) outbreaks caused by the non-native invasive pathogen (*Cronartium ribicola*) (Tomback and Achuff 2010) and native mountain pine beetle (*Dendroctonus ponderosae*) attacks (Logan and Powell 2001, Logan et al. 2010). Decimation of populations in the northern Rocky Mountains has led Canada to list the species as endangered in 2010 ([http://www.cosewic.gc.ca/eng/sct1/searchdetail\\_e.cfm](http://www.cosewic.gc.ca/eng/sct1/searchdetail_e.cfm)). The current and potential loss of this keystone species in the high mountains of California poses serious threats to biodiversity and losses of ecosystem services, since whitebark pine is one of only a few tree species in these settings.

Mountain pine beetles (MPB) are of concern with respect to high elevation conifers and a warming climate. The beetle is a native insect, having co-evolved with western pine forests in fluctuations of periodic disturbance often followed by cleansing fire regime events. More recently, mass beetle infestations have been correlated with increased climatic warming (Mock 2007). Mountain pine beetles require sufficient thermal input to complete the life cycle in one season. Historically, high elevation ecosystems did not meet these conditions. However, due to recent warming trends, there is adequate thermal input at high elevations for the beetle's lifecycle and infestations of whitebark pine are now increasingly common (Logan and Powell 2001). The preponderance of mass infestations at high elevations has been witnessed throughout California—especially in the arid Warner and eastern Sierra Nevada mountains.

In addition to native insects, a non-native fungal pathogen is affecting high elevation forests. In 1910 white pine blister rust (*Cronartium ribicola*) arrived in a British Columbia port and by 1930 had spread to southern Oregon, infect-

ing western white pine (*Pinus monticola*) and sugar pine (*Pinus lambertiana*) (Murray 2005) along the way. The lifecycle completion requires WPBR to utilize *Ribes* spp. as alternate hosts. In late summer, spores from *Cronartium ribicola* are blown from the *Ribes* host and then enter 5-needle pines through stomata. Upon successful entry, hyphae grow, spread through the phloem, then ultimately swell and kill tissue above the site of infection. Infected trees can survive for over 10 years, but the infection inhibits reproduction (Murray 2005). For species like WBP, which live in fringe habitat and therefore delay reproductive events until conditions are optimal, having an infection that further inhibits cone production is a dangerous proposition. The fungus is found on foxtail and whitebark pines in northwest California (Maloy 2001) where variability in microsite infestation occur (Ettl 2007). On Mount Ashland in the Siskiyou Mountains, blister rust has infected 4 of the 9 WBP trees in the population (Murray 2005). All five-needle native western pines have shown some heritable resistance in the past 100 years (Schoettle et al. 2007), but enduring an infection works against a long-lived pine's survival strategy. Populations of whitebark pine did not evolve to withstand fungal infections.

Seedling establishment for organisms that are on the ecological edge, like WBP, is also jeopardized because of the effects of climate change. Causes of unsuccessful seedling recruitment are many but at high elevation include the effects of fire suppression over the past 100 years. While fire has never been a common phenomenon in high-elevation forests, a shift in fire regime occurred in WBP populations during the Holocene, around 4500 years ago. Before that time fire was not a significant factor in WBP ecology but since has become significant (Murray 2005). The introduction of fire regime suppression in the 1930s is another factor in maintaining whitebark populations. The lack of fire, when coupled with effects of climate change, could also lead to population decline. Whitebark pines need open space for seedling establishment and historically some of this open space has been created by fire events. Fire suppression has also led to increased fire severity and intensity which could be compounded by pathogens. If blister rust and mountain pine beetles continue to move into the high elevations of California, they will potentially generate more dead and downed wood. While considering the potential for the risk of stand replacing fire, this would not mimic historical fire regimes—which have been of low intensity and often focused on individual trees by lightning strikes (Murray 2007).

For more images and discussion of whitebark pine forest health in California see supplementary document (Kauffmann 2014).




Figure 1: Range of whitebark pine by Michael Kauffmann

# Introduction

Mapping of whitebark pine occurrence and status/threat has been done primarily using aerial imagery in the National Forests of California by the US Forest Service, including the Pacific Southwest Region - Remote Sensing Lab's CALVEG classification system and maps. The existing USFS vegetation tiles are a result of a 2004-2005 Classification and Assessment with LANDSAT of Visible Ecological Groupings (CALVEG) map product, source imagery ranging from 2002-2009 (USFS 2013c). Even though tile data is continually updated, many stands have not been visited in the field to confirm the accuracy of CALVEG vegetation types. Additionally, little field assessment has been done in the state to identify the presence of whitebark pine, its abundance and status.

The California Native Plant Society (CNPS), working in collaboration with the US Forest Service, initiated field surveys in the summer/fall of 2013 to assess the extent and status of whitebark pine in areas lacking ground surveys in California. Three national forests in the Sierra Nevada and four national forests in the Cascades and Klamath Mountains were selected for field surveys in 2013.

The goals of the field assessments were to verify distribution and status of whitebark pine, ground-truth polygons designated by CALVEG as Whitebark Pine Regional Dominance Type, conduct modified rapid assessments and reconnaissance surveys (recons) on whitebark pine and related stands, and check the USDA Forest Service (USFS) Margins' dataset points for changes in mortality of whitebark pine due to mountain pine beetle and white pine blister rust, if time allowed. Locations within national forests were targeted for the assessment based on potential occurrence of healthy stands in high elevations within the western-most range for the species. Post field assessment, photo interpretation and delineation of whitebark pine extent beyond field surveyed areas were also conducted.

## Methods and Materials

The California Native Plant Society (CNPS) obtained existing GIS data from various sources including the USFS Pacific Southwest - Region Remote Sensing Lab's CALVEG maps (USFS 2013c), USFS Forest Health Technology Enterprise Team's National Insect and Disease Risk Model (USFS 2013a) Host species layers, USFS Pacific Southwest Regional Forest Health and Monitoring Aerial Detection Survey Data (USFS 2013b), USFS Forest Health Protection Margins dataset (Bokach 2013), USFS Forest and Inventory Analysis database (USFS 2013d), The Consortium of California Herbaria (UC Berkeley 2013), USFS Central Sierra Province Ecologist-Becky Estes, USFS Southern Sierra Nevada Province Ecologist - Marc Meyer, National Park Service (NPS) Sierra Nevada Network Inventory and Monitoring Program Ecologist - Jonathan Nesmith, US Geological Survey (USGS) Western Ecological Research Center Ecologist - Nathan Stephenson, California Department of Fish and Wildlife (CDFW) Wildlife Biologist - Pete Figura and USFS Northern California Shared Service Center Entomologist - Cynthia Snyder. In addition, we used older sources of whitebark pine distribution in the state for context (Griffin and Critchfield 1972) and for lone populations or individuals not delineated or attributed by CALVEG (Consortium of California Herbaria, 2014).

CNPS also reviewed existing protocols for evaluating whitebark pine vegetation and insect/disease impacts. These protocols included the NPS Standard Operation Procedures for monitoring White Pine (USDOJ 2012), Whitebark Pine Ecosystem Foundation (Tomback, et al. 2005), Whitebark Pine Inventory and Monitoring Plot protocol (USFS 2013e) and several government research and staff reports (i.e., Millar et al. 2012, Simons and Cluck 2010, Figura 1997, McKinney et al. 2011, and Maloney et al. 2012). We also discussed the existing protocols for assessing whitebark pine vegetation with USFS staff, including Marc Meyer and Shana Gross.

Upon evaluating existing datasets and obtaining input from local National Forest staff, we identified areas to further ground-truth to better determine the distribution and health/status of whitebark pine on the National Forest lands. Priorities included sampling within wilderness lands and identifying areas with low-levels of insect/disease impact.

We selected the CNPS/CDFW Vegetation Rapid Assessment protocol (see Appendix 2) to gather information on occurrence, habitat, and impacts of stands with whitebark pine. We modified this protocol to include signs of Mountain Pine Beetle (MPB) and White Pine Blister Rust (WPBR), and overall whitebark pine status/health. The modified

rapid assessment aimed to gather as much information on whitebark pine health without spending a significant amount of time establishing plots or collecting data on individual trees. Therefore, the survey technique was stand based to assess the extent of whitebark pine vegetation across broad areas in a short amount of time. Sampling included pure stands, mixed conifer stands, and high elevation krummholz, as long as whitebark pine was deemed a component.

The modifications to the rapid assessment included additional information from Marc Meyer's 'Whitebark Pine Inventory and Monitoring Plot Protocol' such as; whitebark pine impacts from MPB and WPBR, MPB level of attack and % of WBP cones (female). Other protocol information added included; # of individual clumps/stems per area, phenology of WBP (% vegetative, % male flowers and % fruiting) and overall site/ occurrence quality/viability (site + population) from the California Natural Diversity Database (CNDDDB). Since MPB attack and WPBR infestation were the main disturbance of interest to be recorded, USFS Pathologists and Entomologists were contacted for visual aids for proper whitebark pine health assessment. Subsequently, comprehensive field guides were made for recognizing symptoms and signs of MPB and WPBR attack (Kauffmann, 2014).

The reconnaissance (recon) form used for the assessment takes pertinent information from the CNPS/CDFW Vegetation Rapid Assessment protocol to gather simplified, general information about a stand (see Appendix 2). Since the goal of the assessment was to gather information on healthy stands of WBP over a large area, the three purposes of the recon form were to collect data either on 1) WBP stands that were largely diseased or infested, 2) stands attributed as WBP by CALVEG but were incorrect, or 3) WBP stands that were close to stands sampled by a Rapid Assessment.

Areas that were selected for sampling in the Klamath National Forest were based on several approaches including identifying and locating populations that were not yet verified, stand accessibility by road, and wilderness settings that were predicted to have been affected by beetles or rust. These assessment areas were also based on places that Michael Kauffmann had already identified as whitebark pine habitat/population centers (Kauffmann 2012, 2013). He has spent extensive time over the past ten years mapping the conifers of the Klamath Mountains, but not the entirety of the Klamath National forest. Therefore, a significant amount of time was spent in the Cascade Mountains north and east of Mount Shasta to find and assess these isolated populations of which little was known. See recommendations section for areas that need future surveying. Deem Burton, with the Happy Camp Ranger District, also provided invaluable information about species distribution before this project began.

## Results

In the Klamath National Forest, I surveyed for and mapped whitebark pine (WBP) in the Goosenest Ranger District and Scott River Ranger District. I found that WBP is inhabiting a variety of ecological niches based on climate, geography, geology, and the synergistic affects of competition from other species. All of these niches, created primarily by elevation, are often "sky islands" that provide small-scale, isolated habitats for WBP across the forest (see maps below). The first habitat type is in the Cascades (east of I-5) where whitebark has been found between 1,980-2,590m (6,500-8,500ft). The upper elevation for the species (outside of the flanks of Mount Shasta) is limited by the height of the peaks themselves. In the Klamath Mountains (west of I-5), the range of elevations is slightly broader at 1,825-2,743m (6,000-9,000ft) but is again limited as the highest peaks, Thompson Peak and Mount Eddy, at 2,750m (9,023ft).

Conifer associates in California's Cascades include white fir (*Abies concolor*), Shasta fir (*Abies magnifica* var. *shastensis*), mountain hemlock (*Tsuga mertensiana*), Lodgepole pine (*Pinus contorta*), western white pine (*Pinus monticola*). In the Klamath Mountains, lodgepole pine are uncommon but others previously listed are common associates. Others include the Klamath endemic foxtail pine (*Pinus balfouriana*) which are nearly synonymous with WBP, and rarely Brewer spruce (*Picea breweriana*), subalpine fir (*Abies lasiocarpa*), Doulgas-fir (*Pseudotsuga menziesii*), and Pacific yew (*Taxus brevifolia*). When present, white and Shasta fir along with mountain hemlock were typically young recruits that appear to be pioneering habitat (encroaching) upon stands of whitebark pine.



### • Goosenest Ranger District - the Cascade Mountains to the north and east of Mount Shasta

This area was a major focus of the summer field work based on the relative shortage of previously collected data. Several permanent plots have been set up in the northern part of this area, including on Ball Mountain and Goosenest, but other data for the southern Goosenest RD was absent. In addition to being quite remote, often only accessible with 4wd, the southern Goosenest is under heavy pressure by the logging industry. A checkerboard of private in-holdings mix with National Forest and logging was often either occurring or had recently occurred within whitebark pine habitat. Often, remnant whitebark pine were left with the completion of logging activities (which took hemlock, western white pine and lodgepole pine instead). There was also salvage logging of lodgepole pine that have been killed by mountain pine beetle in the Whaleback region.

On mountaintops and ridgelines, north-facing slopes were often decimated by mountain pine beetle. These stands were always a mix of lodgepole and whitebark pine with occasional Shasta fir and mountain hemlock. The common pattern is that low species diversity and smaller individual trees are invoked by mesic south-facing slopes with higher species diversity and most vigorous and larger trees inhabiting the mesic north-slopes. MPB are commonly infesting trees on north-slopes because they provide the most concentrated and exploitable resources. This pattern of mortality was found on the Whaleback, the unnamed ridge to the north of Antelope Creek Lakes (private property), as well as the ridgeline between Antelope Creek Lakes and Rainbow Mountain. These areas had mortality reaching and often exceeding 50%. To a lesser degree the Haight Mountain region was exhibiting this pattern but mortality was lower on average. The Ash Creek Butte region appears to be generally unaffected by rust or mountain pine beetles at the time of this writing but this may be due to most of the trees here being smaller and only surviving on the south-facing slopes with north-facing slopes being steep and generally uninhabitable.

### • Scott River Ranger District - the Klamath Mountains

Though time in the Marble Mountains was limited by weather, extensive ground-truthing around Boulder Peak was done. This high table land is one of the most important habitats for WBP in all of the Klamath Mountains. I estimated the overall mortality of five-needle pines to average 5% for whitebark (MPB) and 10% for foxtail pine (MPB). Other enriched montane forest associates between Upper Wright Lake and Boulder Peak include white fir (*Abies concolor*), Shasta fir (*Abies magnifica* var. *shastensis*), subalpine fir (*Abies lasiocarpa*), and mountain hemlock (*Tsuga mertensiana*). These associates do not occur together in this combination anywhere else in the world! This population must be the main feeder for all the other meta-populations isolated on nearby mountaintops and ridgelines across the northern Marble Mountains. Due to the fecundity (at least historically) and extensive size of this population center, Clark's nutcracker were common and must consistently spread seeds to nearby, lower elevation mountain tops, thus sustaining isolated microsites where trees eke out an existence on the last remaining sky island habitats. Box Camp Mountain is a fine example of this phenomenon.

The area around South China and China mountains are mostly on serpentine soils. Because of this, they are already surviving on less than ideal habitat. Unlike other regions of the north state where healthy trees are dying on the north slopes, on the serpentine trees are generally on the south-facing slopes. This habitat supports the most vigorous and healthy trees, outside the competition from other conifers like firs and hemlocks on north slopes. Mortality from MPB and infection from WPBR was found within the small, isolated population on South China Mountain.

### • Salmon River Ranger District

I did not visit the Salmon River Ranger District for this summer's field work. However, whitebark pine occur sparsely on the north face of Thompson as well around Caribou Lake and Caribou Mountain. It would be prudent to develop a better map of the species in the high Trinity Alps.

# Conclusions, Discussion, and Recommendations

The whitebark pine field work in the Klamath National Forest was important in assessing the overall distribution of this vegetation, including significant increases in mapped areas of whitebark pine compared to previous delineations from remotely sensing. The increase in mapped area for the Marble Mountain Wilderness Area was substantial with at least 5 new populations identified and the size of previously identified populations expanded significantly in several cases. While this report is not final in its conclusions and range maps, it is comprehensive as of January 2014.

Using the California Natural Diversity Database (CNDDDB) protocol for documenting overall quality and viability of whitebark pine stands observed in the Marble Mountain and surrounding National Forest areas, we conclude that, overall, populations had good to excellent viability (probability of persistence) over the next 20 years. With the exception of the Cascades north and east of Mount Shasta, whitebark pine in the Klamath NF are relatively healthy.

## • Goosenest Ranger District

1. Ground-truth Willow Creek Mountain and Garner Mountain
2. Look close at the West Haight Mountain stand and possibly pursue the designation of a botanical area
3. Set up long term monitoring in the Antelope Creek Research Natural Area because it is one of the most susceptible to decline due to the narrow ecological amplitude offered here.

## • Scott River

1. Ground truth the Heather Lake Pluton, between Man Eaten and Cliff lakes and the “Big Ridge” between Black Marble Mountain and King’s Castle in the Marble Mountain Wilderness.
2. Set up long term monitoring plots in the Boulder Peak Region. This is one of the most extensive stands of whitebark pine in the Klamath Mountains and most likely serves as a “feeder” population for the smaller mountain-top stands nearby. This area is critical to the future of whitebark in the Klamath Mountain, in my opinion.
3. Ground truth and create a better map for the Russian Wilderness area, especially the higher peaks around Upper Albert and Big Blue lakes. I’m actually not 100% the trees are here, but they should be!
4. Assess the population on China Mountain and set up a permanent plot on South China Mountain.
5. Work with Shasta-Trinity to ground-truth the Cory Peak Botanical and Geological area to verify or nullify species occurrence.

## • Salmon River Ranger District

1. Work with the Weaverville and Big Bar RDs on the Shasta-Trinity to map and ground-truth the extent of WBP along the Stuarts Fork-Salmon Divide (this is some steep country!).
2. I believe that WBP could occur in the Dorleska Mine region in the Big Flat Region. This area should be ground-truthed and mapped

## • Across the Klamath National Forest

1. Create a map to target areas where encroachment from firs and hemlocks is an issue and consider managing for this problem

Lastly, this report is not comprehensive; it was based upon the available funding, resources and USDA Forest Service staff schedules in 2013. The draft map of whitebark pine distribution (see Figure 2) is therefore not complete but hopefully provides an updated version of whitebark pine distribution from field surveys and aerial interpretation with limited modeled data. The modeled data that is presented from CALVEG in Figure 2 is used to provide areas of data gaps where future field assessments are needed.

## More resources for whitebark pine in northern California:

Keeler-Wolf, Todd. 1990. Ecological surveys of FS research natural areas in California.  
[http://www.fs.fed.us/psw/publications/documents/psw\\_gtr125/](http://www.fs.fed.us/psw/publications/documents/psw_gtr125/)

- o Crater Creek RNA, Klamath
- o Mt. Eddy RNA, Shasta Trinity
- o Sugar Creek RNA, Klamath
- o Antelope Creek Lakes, Klamath

Table 1. Area of whitebark pine populations by national forest region in northern California

Forest	Region	Acres	Hectares
Klamath National Forest	South Goosenest (5 polygons with a small amount in the Shasta-Trinity NF)	2,631	1,065
	North Goosenest (3 polygons)	152	62
	Marble Mountain Wilderness	4,721	1,911
	Russian Wilderness	630	255
	China Mountain Region (some in Shasta-Trinity)	609	246
	<b>total</b>	<b>9,198</b>	<b>3,722</b>
Shasta-Trinity National Forest	Mount Eddy Region	6,048	2,448
	Mount Shasta	11,595	4,692
	Trinity Alps Wilderness	5,671	2,295
	Also see China Mountain and South Goosenest above		
	<b>total</b>	<b>22,039</b>	<b>8,919</b>
Lassen National Forest (including Lassen N.P.)	Within Lassen National Park	11,435	4,628
	Thousand Lakes Wilderness	645	261
	Burney Mountain	15	6
	<b>total</b>	<b>12,095</b>	<b>4,895</b>
Modoc National Forest	Buck Mountain Region	2,401	826
	South Warner Wilderness	20,125	8,548
	Middle Warners	448	181
	North Warners	3,884	1572
	<b>total</b>	<b>26,858</b>	<b>11,127</b>
<b>Total acreage in the four forest regions of Northern California</b>		<b>70,906</b>	<b>28,633</b>


Table 2. Rapid Assessment summary, Klamath NF (note that Ash Creek Butte site falls within Shasta-Trinity NF boundaries).

DbaseID	County	Ranger District	Wilderness	Site name	Alliance	Estimated Pct Cover PIAL	Altitude (m)	Impacts
WBP0027	Siskiyou	Goosenest		Antelope Creek	Western North American Montane Sclerophyll Scrub Group	2	2043	
WBP0028	Siskiyou	Goosenest		Antelope Creek	Pinus albicaulis	15	2186	MPB (50%), Rust (31%)
WBP0037	Siskiyou	Goosenest		Haight	Tsuga mertensiana	4	2146	Logging (medium), MPB (25%), Rust (50%)
WBP0038	Siskiyou	Goosenest		Haight	Abies magnifica - Abies concolor	5	2138	MPB (80%)
WBP0039	Siskiyou	Goosenest		Haight	Abies concolor	9	2185	Rust (25%)
WBP0040	Siskiyou	Goosenest		Haight	Abies magnifica - Abies concolor	5	2241	MPB (90%), Rust (50%)
WBP0041	Siskiyou	Goosenest		Haight	Pinus albicaulis	30	2414	Rust (15%)
WBP0042	Siskiyou	Goosenest		Haight	Pinus albicaulis	8	2278	MPB (35%)
WBP0065	Siskiyou	Goosenest		Whaleback	Pinus contorta subsp. murrayana	8	2431	MPB (40%), Rust (40%)
WBP0066	Siskiyou	Goosenest		Whaleback	Pinus albicaulis	20	2579	MPB (5%), Rust (20%)
WBP0067	Siskiyou	Goosenest		Whaleback	Pinus albicaulis	15	2349	Rust (20%)
WBP0068	Siskiyou	Goosenest		Whaleback	Pinus contorta subsp. murrayana	10	2298	Logging (high)
WBP0029	Siskiyou	McCloud		Ash Creek Butte	Pinus albicaulis	10	2320	MPB (10%), Rust (10%)
WBP0030	Siskiyou	McCloud		Ash Creek Butte	Pinus albicaulis	23	2520	MPB (10%), Rust (5%)
WBP0031	Siskiyou	McCloud		Ash Creek Butte	Pinus albicaulis	25	2452	MPB (5%)
WBP0054	Siskiyou	Scott River		South China Mtn	Pinus balfouriana	5	2457	MPB (70%), Rust (50%)
WBP0055	Siskiyou	Scott River		South China Mtn	Pinus balfouriana	5	2395	Rust (10%)
WBP0036	Siskiyou	Scott River	Marble Mountain	Boulder Peak	Abies concolor	5	1985	Grazing (low), MPB (50%), Rust (30%)

Table 3. *Pinus albicaulis* attributes from Rapid Assessments in Klamath NF (note that Ash Creek Butte site falls within Shasta-Trinity NF boundaries). N/A is used when data was not recorded.

<b>DbaseID</b>	<b>Site name</b>	<b>Clumps per hectare</b>	<b>Percent Vegetative</b>	<b>Percent Flowering</b>	<b>Percent Fruiting</b>	<b>Mortality by MPB</b>	<b>Total Mortality</b>	<b>Quality</b>
WBP0027	Antelope Creek	5.0	100			0	0	Excellent
WBP0028	Antelope Creek	15.0	92		8	50%	50%	Fair
WBP0037	Haight	20.0	90	10		25%	25%	Poor
WBP0038	Haight	10.0	100			90%	90%	Poor
WBP0039	Haight	12.5	90		10	0	10%	Good
WBP0040	Haight	10.0	20		80	90%	90%	Poor
WBP0041	Haight	25.0	20		80	0	0	Excellent
WBP0042	Haight	30.0	70	30		30%	30%	Good
WBP0065	Whaleback	20.0	100			40%	40%	Fair
WBP0066	Whaleback	25.0	80		20	0	0	Fair
WBP0067	Whaleback	15.0	90			0	0	Excellent
WBP0068	Whaleback	n/a	n/a	n/a	n/a	n/a	0	Good
WBP0029	Ash Creek Butte	15.0	90		10	10%	10%	Good
WBP0030	Ash Creek Butte	25.0	80	20		10%	10%	Good
WBP0031	Ash Creek Butte	50.0	60		40	5%	5%	Excellent
WBP0054	South China Mtn	20.0	100			70%	70%	Poor
WBP0055	South China Mtn	10.0	100			0	0	Excellent
WBP0036	Boulder Peak	7.5	70	20	20	30%	30%	Fair

Figure 2: Overview Maps of 2013 Locations Visited on the National Forest






Figure 3: Goosenest Ranger District - Detailed Maps




Figure 4: Goosenest Ranger District - North







Figure 5: Goosenest Ranger District - South

Figure 6: Scott River Ranger District - Overview






Figure 7: Scott River Ranger District - China Mountain Region




Figure 8: Scott River Ranger District - Marble Mountain Overview




Figure 9: Scott River Ranger District - Marble Mountain Overview




Figure 10: Scott River Ranger District - Russian Wilderness Overview




Figure 11: Range of whitebark pine in the divide between the Big Bar, Weaverville, and Salmon River Ranger Districts - Trinity Alps Wilderness

# Goosenest Ranger District - Klamath National Forest

## The Whaleback



*Figure 12: Looking south through a whitebark pine from the Whaleback*



*Figure 13: Looking east to an unnamed peak. Notice the concentration of mortality of lodgepole and whitebark pine on the north-facing slope*



*Figure 14: High mortality due to mountain pine beetle*



# Gooseneck Ranger District - Klamath National Forest

## Haight Mountain Region



*Figure 15: Extensive stand of whitebark pine spreads through the basin between Haight Mountain (back left) and West Haight*



*Figure 16: White fir are encroaching upon stands of whitebark pine along south-facing ridgelines near West Haight Mountain.*

*Figure 17: The view east from the summit of West Haight Mountain, through whitebark pine toward the Medicine Lake Highlands*



## McCloud Ranger District - Shasta-Trinity National Forest Ash Creek Butte



*Figure 18: From the summit of Ash Creek Butte, looking west toward Mount Shasta*

## Goosenest Ranger District - Klamath National Forest Antelope Creek RNA



*Figure 19: From the steep ridgelines of the Antelope Creek Research Natural Area, mountain pine beetle kill was common (~50% of trees). Here looking west to Ash Creek Butte and Mount Shasta.*

# Scott River Ranger District - Klamath National Forest

## Boulder Peak Regions - Marble Mountain Wilderness



*Figure 20: Whitebark pine mix with foxtail pine (*Pinus balfouriana*) and mountain hemlock (*Tsuga mertensiana*) along the high elevation tablelands adjacent to Boulder peak in the Marble Mountain Wilderness*



*Figure 21: The view to the west from Boulder peak, with whitebark pines framing the wilderness' namesake, Marble Mountain*

## Scott River Ranger District - Klamath National Forest Boulder Peak Regions - Marble Mountain Wilderness



*Figure 22: Lower Wright Lake Basin holds a sparse mixture of whitebark and foxtail pines.*



*Figure 23: An enriched montane forest near Upper Wright Lake holds numerous conifer species including whitebark pine, foxtail pine (*Pinus balfouriana*), white fir (*Abies concolor*), Shasta fir (*Abies x shastensis*), subalpine fir (*Abies lasiocarpa*), and mountain hemlock (*Tsuga mertensiana*). These associations do not occur anywhere else in the world!*

# Scott River Ranger District - Klamath National Forest

## South China Mountain



*Figure 24: Where serpentine meets gabbro, whitebark pine survive only in the highest elevations of South China Mountain*



*Figure 25: Branch flagging from white pine blister rust is evident on the whitebark pine (left) and mountain pine beetle completely killed foxtail pines (right)*

## Literature Cited

- Arno, S.F., R.J. Hoff. 1989. Silvics of Whitebark Pine (*Pinus Albicaulis*). USDA Forest Service Technical Report. INT-253.
- BLM. 2014. Federal and State managed lands in California and portions northwest Nevada. Bureau of Land Management, California State Office, Mapping Sciences 5/15/2009. Data available at: <http://www.blm.gov/ca/gis/>
- Bokach, M.. 2013. Margin's dataset. USDA Forest Service, Forest Health Protection Program.
- Bunn, A.G., L.J. Graumlich, D.L Urban. 2005. Trends in twentieth-century tree growth at high elevations in the Sierra Nevada and White Mountains, USA. *The Holocene* 15:481-488.
- CNDDDB. 2014. California Natural Diversity Database (CNDDDB). California Department of Fish and Game, Biogeographic Data Branch, Vegetation Classification and Mapping Program, Sacramento, CA. <http://www.dfg.ca.gov/biogeodata/cnddb/>
- Charlet, D.A. and B. Keimel. 1996 (2012). Atlas of Nevada conifers: a phytogeographic reference. University of Nevada Press, Reno, Nevada. Also, personal communication with author (2012).
- Ettl, G. J. 2007. Ecology of Whitebark Pine in the Pacific Northwest. Proceedings of the Conference Whitebark Pine: A Pacific Coast Perspective USDA Forest Service p20-22
- Figura, P. J. 1997. Structure and dynamics of whitebark pine forests in the South Warner Wilderness, northeastern California. A thesis presented to the faculty of Humboldt State University, Humbolt, CA.
- Greater Yellowstone Whitebark Pine Monitoring Working Group. 2007. Interagency Whitebark Pine Monitoring Protocol for the Greater Yellowstone Ecosystem, Version 1.00. Greater Yellowstone Coordinating Committee, Bozeman, MT <http://www.greateryellowstonescience.org/subproducts/14/72>
- Griffin, J. R., and W.B. Critchfield. 1976. The Distribution of Forest Trees in California. USDA Forest Service: Berkeley, CA.
- Kane, J.M. et al. 2013. Resin duct characteristics associated with tree resistance to bark beetles across lodgepole and limber pines. *Oecologia*. December 2013. <http://dx.doi.org/10.1007/s00442-013-2841-2>
- Kauffmann, M. 2012. Conifer Country: A natural history and hiking guide to the 35 conifers of the Klamath Mountain region. Backcountry Press, Kneeland, California.
- Kauffmann, M. 2013. Conifers of the Pacific Slope. Backcountry Press, Kneeland, California.
- Kauffmann, M. 2014. Whitebark Pine Forest Health in California. Unpublished report. California Native Plant Society Vegetation Program, Sacramento, CA. [<http://pacslope-conifers.com/conifers/pine/wbp/CNPS-Reports/WBP-Forest-Health-California.pdf>]
- Keeler-Wolf, Todd. 1990. Ecological surveys of FS research natural areas in California. Available at: [http://www.fs.fed.us/psw/publications/documents/psw\\_gtr125/](http://www.fs.fed.us/psw/publications/documents/psw_gtr125/)
- Lanner, R.M. 1996. Made for each other: A symbiosis of birds and pines. Oxford University Press. New York.
- Little, E.L., Jr. 1971. Atlas of United States trees, volume 1, conifers and important hardwoods: U.S. Department of Agriculture Miscellaneous Publication 1146, 9 p., 200 maps.
- Logan, J.A. and J.A. Powell. 2001. Ghost forests, global warming, and the mountain pine beetle (Coleoptera: Scolytidae). *American Entomologist* (Fall) 47: 160 - 172.
- Logan, J.A., W.W. Macfarlane, and L. Willcox, 2010. Whitebark pine vulnerability to climate-driven mountain pine beetle disturbance in the Greater Yellowstone Ecosystem. *Ecol. Appl.* 20(4): 895–902.
- Maloney, P. E., D.R. Vogler, C.E. Jensen, A.D. Mix. 2012. Ecology of whitebark pine populations in relation to white pine blister rust infection in subalpine forests of the Lake Tahoe Basin, USA: Implications for restoration. *Forest Ecology and Management* 280 (2012) 166–175.
- Maloy, O. C. 2001. White pine blister rust. Online. Plant Health Progress doi:10.1094/PHP-2001-0924-01-HM
- McKinney, S. T., T. Rodhouse, L. Chow, P. Latham, D. Sarr, L. Garrett, L. Mutch. 2011. Long-Term Monitoring of High-Elevation White Pine Communities in Pacific West Region National Parks. USDA Forest Service Proceedings RMRS-P-63.

- Millar, C.I., R. D. Westfall, D. L. Delany, M. J. Bokach. 2012. Forest mortality in high-elevation whitebark pine (*Pinus albicaulis*) forests of eastern California, USA; influence of environmental context, bark beetles, climatic water deficit, and warming. *Canadian Journal of Forest Research* 42: 749–765
- Mock, K.E., et. al. 2007. Landscape-scale genetic variation in a forest outbreak species, the mountain pine beetle (*Dendroctonus ponderosae*). *Molecular Ecology* 16:553-568.
- Murray, M.P. 2005. Our threatened timberlines: the plight of whitebark pine ecosystems. *Kalmiopsis*. 12:25-29
- Murray, M.P. 2007 Fire and Pacific Coast Whitebark Pine. Proceedings of the Conference Whitebark Pine: A Pacific Coast Perspective USDA Forest Service p51-60.
- National Park Service (NPS). 2012. Monitoring White Pine (*Pinus albicaulis*, *P. balfouriana*, *P. flexilis*) Community Dynamics in the Pacific West Region - Klamath, Sierra Nevada, and Upper Columbia Basin Networks, Standard Operating Procedures Version 1.0. Natural Resource Report NPS/PWR/NRR—2012/533.
- Sawyer, J.O. 2004. Conifers of the Klamath Mountains. *Vegetation Ecology, Proceedings of the second conference on Klamath-Siskiyou ecology*. 128–135
- Sawyer, J. O. 2006. Northwest California. University of California Press, Berkeley, CA.
- Schoettle, A.W. and R.A. Sniezko. 2007. Proactive intervention to sustain high-elevation pine ecosystems threatened by white pine blister rust. *Journal of Forestry Research*. 12: 327-336.
- Simons, R., D. Cluck. 2010. Whitebark pine monitoring plot protocol for the Warner Mountains, Modoc National Forest. USDA Forest Service, Forest Health Protection and Modoc National Forest.
- Tomback, D.F., and P. Achuff, 2010. Blister rust and western forest biodiversity: ecology, values and outlook for white pines. *Forest Pathology* 40: 186–225.
- Tomback, D. F., R.E. Keane, W. W. McCaughey, C. Smith. 2004. Methods for Surveying and Monitoring Whitebark Pine for Blister Rust Infection and Damage. Whitebark Pine Ecosystem Foundation, Missoula, MT.
- UC Berkeley. 2013. Consortium of California Herbaria. Data provided by the participants of the Consortium of California Herbaria ([ucjeps.berkeley.edu/consortium/](http://ucjeps.berkeley.edu/consortium/))
- US. Department of the Interior. 2012. Monitoring White Pine (*Pinus albicaulis*, *P. balfouriana*, *P. flexilis*) Community Dynamics in the Pacific West Region - Klamath, Sierra Nevada, and Upper Columbia Basin Networks, Standard Operating Procedures Version 1.0
- USDA Forest Service. 2013a. Host species layers. U.S. Forest Service Forest Health Technology Enterprise Team; 2013 National Insect and Disease Risk Model. Data available at: <http://www.fs.fed.us/foresthealth/technology/nidrm.shtml>
- USDA Forest Service. 2013b. U.S. Forest Service Pacific Southwest Regional Forest Health Monitoring Aerial Detection Survey Data. Data available at: <http://www.fs.usda.gov/detail/r5/landmanagement/resourcemanagement/?cid=stelprdb5347192>
- USDA Forest Service. 2013c. Vegetation mapping. Existing vegetation (evveg) layers. Pacific Southwest Region Remote Sensing Lab. Data available at <http://www.fs.usda.gov/detail/r5/landmanagement/resourcemanagement/?cid=stelprdb5347192>
- USDA Forest Service. 2013d. Forest and Inventory Analysis database. Forest Inventory and Analysis National Program. Data available at: <http://www.fia.fs.fed.us/tools-data/>
- USDA Forest Service. 2013e. Whitebark Pine Inventory and Monitoring Plot Protocol. USFS Region 5 Ecology Program and Forest Health Protection Program.

# Appendix 1: Inventory and Monitoring Protocols and Field Forms from 2013

## CALIFORNIA NATIVE PLANT SOCIETY / DEPARTMENT OF FISH AND GAME PROTOCOL FOR COMBINED VEGETATION RAPID ASSESSMENT AND RELEVÉ SAMPLING FIELD FORM

(Modified for WBP)

July 8, 2013

### Introduction

This protocol describes the methodology for both the relevé and rapid assessment vegetation sampling techniques as recorded in the combined relevé and rapid assessment field survey form dated June 28, 2013. The same environmental data are collected for both techniques. However, the relevé sample is plot-based, with each species in the plot and its cover being recorded. The rapid assessment sample is based not on a plot but on the entire stand, with 12-20 of the dominant or characteristic species and their cover values recorded. For more background on the relevé and rapid assessment sampling methods, see the relevé and rapid assessment protocols at [www.cnps.org](http://www.cnps.org).

### Selecting stands to sample:

To start either the relevé or rapid assessment method, a stand of vegetation needs to be defined.

A stand is the basic physical unit of vegetation in a landscape. It has no set size. Some vegetation stands are very small, such as alpine meadow or tundra types, and some may be several square kilometers in size, such as desert or forest types.

A stand is defined by two main unifying characteristics:

- 1) It has compositional integrity. Throughout the site, the combination of species is similar. The stand is differentiated from adjacent stands by a discernable boundary that may be abrupt or indistinct.
- 2) It has structural integrity. It has a similar history or environmental setting that affords relatively similar horizontal and vertical spacing of plant species. For example, a hillside forest originally dominated by the same species that burned on the upper part of the slopes, but not the lower, would be divided into two stands. Likewise, sparse woodland occupying a slope with very shallow rocky soils would be considered a different stand from an adjacent slope with deeper, moister soil and a denser woodland or forest of the same species.

The structural and compositional features of a stand are often combined into a term called homogeneity. For an area of vegetated ground to meet the requirements of a stand, it must be homogeneous (uniform in structure and composition throughout).

Stands to be sampled may be selected by evaluation prior to a site visit (e.g., delineated from aerial photos or satellite images), or they may be selected on site during reconnaissance (to determine extent and boundaries, location of other similar stands, etc.).

Depending on the project goals, you may want to select just one or a few representative stands of each homogeneous vegetation type for sampling (e.g., for developing a classification for a vegetation mapping project), or you may want to sample all of them (e.g., to define a rare vegetation type and/or compare site quality between the few remaining stands).

For the rapid assessment method, you will collect data based on the entire stand.



## Selecting a plot to sample within in a stand (for relevés only):

Because many stands are large, it may be difficult to summarize the species composition, cover, and structure of an entire stand. We are also usually trying to capture the most information as efficiently as possible. Thus, we are typically forced to select a representative portion to sample.

When sampling a vegetation stand, the main point to remember is to select a sample that, in as many ways possible, is representative of that stand. This means that you are not randomly selecting a plot; on the contrary, you are actively using your own best judgment to find a representative example of the stand.

Selecting a plot requires that you see enough of the stand you are sampling to feel comfortable in choosing a representative plot location. Take a brief walk through the stand and look for variations in species composition and in stand structure. In many cases in hilly or mountainous terrain look for a vantage point from which you can get a representative view of the whole stand. Variations in vegetation that are repeated throughout the stand should be included in your plot. Once you assess the variation within the stand, attempt to find an area that captures the stand's common species composition and structural condition to sample.

## Plot Size

All relevés of the same type of vegetation to be analyzed in a study need to be the same size. Plot shape and size are somewhat dependent on the type of vegetation under study. Therefore, general guidelines for plot sizes of tree-, shrub-, and herbaceous communities have been established. Sufficient work has been done in temperate vegetation to be confident the following conventions will capture species richness:

Herbaceous communities: 100 sq. m plot

Special herbaceous communities, such as vernal pools, fens: 10 sq m plot

Shrublands and Riparian forest/woodlands: 400 sq. m plot

Open desert and other shrublands with widely dispersed but regularly occurring woody species: 1000 sq. m plot

Upland Forest and woodland communities: 1000 sq. m plot

## Plot Shape

A relevé has no fixed shape, though plot shape should reflect the character of the stand. If the stand is about the same size as a relevé, the plot boundaries may be similar to that of the entire stand. If we are sampling streamside riparian or other linear communities, our plot dimensions should not go beyond the community's natural ecological boundaries. Thus, a relatively long, narrow plot capturing the vegetation within the stand, but not outside it would be appropriate. Species present along the edges of the plot that are clearly part of the adjacent stand should be excluded.

If we are sampling broad homogeneous stands, we would most likely choose a shape such as a circle (which has the advantage of the edges being equidistant to the center point) or a square (which can be quickly laid out using perpendicular tapes).

## Definitions of fields in the protocol

**Relevé or Rapid Assessment:** Circle the method that you are using.

### I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION

**Polygon/Stand #:** Number assigned either in the field or in the office prior to sampling. It is usually denoted with a four-letter abbreviation of the sampling location and then a four-number sequential number of that locale (e.g. CARR0001 for Carrizo sample #1). The maximum number of letters/numbers is eight.

**Air photo #:** The number given to the aerial photo in a vegetation-mapping project, for which photo interpreters have already done photo interpretation and delineations of polygons. If the sample site has not been photo-interpreted, leave blank.

**Date:** Date of the sampling.

**Name(s) of surveyors:** The full names of each person assisting should be provided for the first field form for the day. On successive forms, initials of each person assisting can be recorded. Please note: The person recording the data on the form should circle their name/initials.

**GPS waypoint #:** The waypoint number assigned by a Global Positioning System (GPS) unit when marking and storing a waypoint for the sample location. Stored points should be downloaded in the office to serve as a check on the written points and to enter into a GIS.

For relevé plots, take the waypoint in the southwest corner of the plot or in the center of a circular plot.

**GPS name:** The name/number assigned to each GPS unit. This can be the serial number if another number is not assigned.

**Datum:** (NAD 83) The standard GPS datum used is NAD 83. If you are using a different datum, note it here.

**Bearing, left axis at SW pt (note in degrees) of Long or Short side:** For square or rectangular plots: from the SW corner (= the GPS point location), looking towards the plot, record the bearing of the axis to your left. If the plot is a rectangle, indicate whether the left side of the plot is the long or short side of the rectangle by circling "long" or "short" side (no need to circle anything for circular or square plots). If there are no stand constraints, you would choose a circular or square plot and straight-sided plots should be set up with boundaries running in the cardinal directions. If you choose a rectangular plot that is not constrained by the stand dimensions, the short side should run from east to west, while the long side should run from north to south.

**UTM coordinates:** Easting (UTME) and northing (UTMN) location coordinates using the Universal Transverse Mercator (UTM) grid. Record in writing the information from a GPS unit or a USGS topographic map.

**UTM zone:** Universal Transverse Mercator zone. Zone 10 is for California west of the 120th longitude, zone 11 is for California east of 120th longitude, which is the same as the straight portion of California's eastern boundary.

**Error ±:** The accuracy of the GPS location, when taking the UTM field reading. Please record the error units by circling feet (ft), meters (m), or positional dilution of precision (pdop). If your GPS does not determine error, insert N/A in this field.

**Is GPS within stand?** Yes / No Circle “Yes” to denote that the GPS waypoint was taken directly within or at the edge of the stand being assessed for a rapid assessment, or circle “No” if the waypoint was taken at a distance from the stand (such as with a binocular view of the stand).

**If No, cite from waypoint to stand, distance (note in meters) & bearing (note in degrees):** An estimate of the number of meters and the compass bearing from the GPS waypoint to the stand.

**Elevation:** Recorded from the GPS unit or USGS topographic map. Please circle feet (ft) or meters (m).

**Photograph #s:** Write the name or initials of the camera owner, JPG/frame number, and direction of photos (note the roll number if using film). Take four photos in the main cardinal directions (N, E, S, W) clockwise from the north, from the GPS location. If additional photos are taken in other directions, please note this information on the form. Also include overview photos of Whitebark pine.

**Stand Size:** Estimate the size of the entire stand in which the sample is taken. As a measure, one acre is about 4000 square meters (approximately 64 x 64 m), or 208 feet by 208 feet. One acre is similar in size to a football field.

**Plot Size:** If this is a relevé, circle the size of the plot.

**Plot Shape:** Record the length and width of the plot and circle measurement units (i.e., ft or m). If it is a circular plot, enter radius (or just put a check mark in the space).

**Exposure:** (Enter actual ° and circle general category): With your back to the general uphill direction of the slope (i.e., by facing downhill of the slope), read degrees of the compass for the aspect or the direction you are standing, using degrees from north, adjusted for declination. Average the reading over the entire stand, even if you are sampling a relevé plot, since your plot is representative of the stand. If estimating the exposure, write “N/A” for the actual degrees, and circle the general category chosen. “Variable” may be selected if the same, homogenous stand of vegetation occurs across a varied range of slope exposures. Select “all” if stand is on top of a knoll that slopes in all directions or if the same, homogenous stand of vegetation occurs across all ranges of slope.

**Steepness:** (Enter actual ° and circle general category): Read degree slope from a compass or clinometer. If estimating, write “N/A” for the actual degrees, and circle the general category chosen.. Make sure to average the reading across the entire stand even if you are sampling in a relevé plot.

**Topography:** First assess the broad (Macro) topographic feature or general position of the stand in the surrounding watershed, that is, the stand is at the top, upper (1/3 of slope), middle (1/3 of slope), lower (1/3 of slope), or bottom. Circle all of the positions that apply for macrotopography.

Then assess the local (Micro) topographic features or the lay of the area (e.g., surface is flat or concave). Circle only one of the microtopographic descriptors.

**Geology:** Geological parent material of site. If exact type is unknown, use a more general category (e.g., igneous, metamorphic, sedimentary). See code list for types.

**Soil Texture:** Record soil texture that is characteristic of the site (e.g., coarse loamy sand, sandy clay loam). See soil texture key and code list for types.

**Upland or Wetland/Riparian (circle one):** Indicate if the stand is in an upland or a wetland. There are only two options. Wetland and riparian are one category. Note that a site need not be officially delineated as a wetland to qualify as such in this context (e.g., seasonally wet meadow).

**% Surface cover (abiotic substrates).** It is helpful to imagine “mowing off” all of the live vegetation at the base of the plants and removing it – you will be estimating what is left covering the surface. The total should sum to 100%. Note that non-vascular cover (lichens, mosses, cryptobiotic crusts) is not estimated in this section.

**% Water:** Estimate the percent surface cover of running or standing water, ignoring the substrate below the water.

**% BA Stems:** Percent surface cover of the plant basal area, i.e., the basal area of stems at the ground surface. Note that for most vegetation types BA is 1-3% cover. Estimate for a set area (e.g., 400 m<sup>2</sup>) of BA to help calibrate on this % (on average % is between 1.5-4.5% for conifers)

**% Litter: Percent surface cover of litter, duff, or wood on the ground.**

**% Bedrock:** Percent surface cover of bedrock.

**% Boulders:** Percent surface cover of rocks > 60 cm in diameter.

**% Stone:** Percent surface cover of rocks 25-60 cm in diameter.

**% Cobble:** Percent surface cover of rocks 7.5 to 25 cm in diameter.

**% Gravel:** Percent surface cover of rocks 2 mm to 7.5 cm in diameter.

**% Fines:** Percent surface cover of bare ground and fine sediment (e.g. dirt) < 2 mm in diameter.

**% Current year bioturbation:** Estimate the percent of the sample or stand exhibiting soil disturbance by fossorial organisms (any organism that lives underground). Do not include disturbance by ungulates. Note that this is a separate estimation from surface cover.

**Past bioturbation present?** Circle Yes if there is evidence of bioturbation from previous years.

**% Hoof punch:** Note the percent of the sample or stand surface that has been punched down by hooves (cattle or native grazers) in wet soil.

**Fire Evidence:** Circle Yes if there is visible evidence of fire, and note the type of evidence in the “Site history, stand age and comments section,” for example, “charred dead stems of *Quercus berberidifolia* extending 2 feet above resprouting shrubs.” If you are certain of the year of the fire, put this in the Site history section.

**Site history, stand age, and comments:** Briefly describe the stand age/seral stage, disturbance history, nature and extent of land use, and other site environmental and vegetation factors. Examples of disturbance history: fire, landslides, avalanching, drought, flood, animal burrowing, or pest outbreak. Also, try to estimate year or frequency of disturbance. Examples of land use: grazing, timber harvest, or mining. Examples of other site factors: exposed rocks, soil with fine-textured sediments, high litter/duff build-up, multi-storied vegetation structure, or other stand dynamics.

**Disturbance code / Intensity (L,M,H):** List codes for potential or existing impacts on the stability of the plant community. Characterize each impact each as L (=Light), M (=Moderate), or H (=Heavy). For invasive exotics, divide the total exotic cover (e.g. 25% *Bromus diandrus* + 8% *Bromus madritensis* + 5% *Centaurea melitensis* = 38% total exotics) by the total % cover of all the layers when added up (e.g. 15% tree + 5% low tree + 25% shrub + 40% herbs = 85% total) and multiply by 100 to get the % relative cover of exotics (e.g. 38% total exotics/85% total cover = 45% relative exotic cover). L = 0-33% relative cover of exotics; M = 34-66% relative cover, and H = > 66% relative cover. See code list for impacts.

List percent of WBP impacted by Mountain Pine Beetle (39-MPB/L/approx. % impacted) and White Pine Blister Rust (40-WPBR/H/approx. % impacted) within the stand. For Mountain Pine Beetle, search the bole for entry holes (reddish colored pitch) or frass. For WPBR, search for ‘signs’ of an active canker (i.e., a canker with visible aecia, or fruiting bodies containing spores), or ‘symptoms’ of any of the following five indicators: rodent chewing, flagging, swelling, roughened bark, and oozing sap. Explain signs and symptoms in the notes and take photos when necessary.

## II. HABITAT AND VEGETATION DESCRIPTION

### California Wildlife-Habitat Relationships (CWHR)

For CWHR, identify the size/height class of the stand using the following tree, shrub, and/or herbaceous categories. These categories are based on functional life forms.

**Tree DBH:** Circle one of the tree size classes provided when the tree canopy closure exceeds 10 percent of the total cover, or if young tree density indicates imminent tree dominance. Size class is based on the average diameter at breast height (dbh) of each trunk (standard breast height is 4.5ft or 137cm). When marking the main size class, make sure to estimate the mean diameter of all trees over the entire stand, and weight the mean if there are some larger tree dbh's. The "T6 multi-layered" dbh size class contains a multi-layered tree canopy (with a size class T3 and/or T4 layer growing under a T5 layer and a distinct height separation between the classes) exceeding 60% total cover. Stands in the T6 class need also to contain at least 10% cover of size class 5 (>24" dbh) trees growing over a distinct layer with at least 10% combined cover of trees in size classes 3 or 4 (>11-24" dbh). This is weighted: In your representative area add number of trees for each category and record above (T1,T2,T3, etc). Can square root later to get the weighted average for this category (if there are many sizes).

**Shrub:** Circle one of the shrub size classes provided when shrub canopy closure exceeds 10 percent (except in desert types) by recording which class is predominant in the survey. Shrub size class is based on the average amount of crown decadence (dead standing vegetation on live shrubs when looking across the crowns of the shrubs).

**Herb:** Circle one of the herb height classes when herbaceous cover exceeds 2 percent by recording the predominant class in the survey. Note: This height class is based on the average plant height at maturity, not necessarily at the time of observation.

**Desert Palm/Joshua Tree:** Circle one of the palm or Joshua tree size classes by averaging all the stem-base diameters (i.e. mean diameter of all stem-base sizes). Diameter is measured at the plant's base above the bulge near the ground.

**Desert Riparian Tree/Shrub:** Circle one of the size classes by measuring mean stem height (whether tree and/or shrub stand).

### Overall Cover of Vegetation

Provide an estimate of cover for the following categories below (based on functional life forms). Record a specific number for the total aerial cover or "bird's-eye view" looking from above for each category, estimating cover for the living plants only. Litter/duff should not be included in these estimates. The porosity of the vegetation should be taken into consideration when estimating percent cover (how much of the sky can you see when you are standing under the canopy of a tree, or how much light passes through the canopy of the shrub layer?).

To come up with a specific number estimate for percent cover, first use generalized cover classes as reference aids such as the CWHR cover classes (<2%, 2-9%, 10-24%, 25-39%, 40-59%, 60-100%) or the modified Braun-Blanquet cover-abundance scale (<1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, >75%). While keeping these intervals in mind, you can then refine your estimate to a specific percentage for each category below.

**% Total NonVasc cover:** The total cover of all lichens, bryophytes (mosses, liverworts, hornworts), and cryptogammic crust on substrate surfaces including downed logs, rocks and soil, but not on standing or inclined trees or vertical rock surfaces.

**% Total Vasc Veg cover:** The total cover of all vascular vegetation taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute vegetation cover, disregarding overlap of the various tree, shrub, and/or herbaceous layers and species. Could use densitometer to calibrate, but sometimes this provides an over-estimate.

### **% Cover by Layer**

**% Conifer Tree /Hardwood Tree:** The total foliar cover (considering porosity) of all live tree species, disregarding overlap of individual trees. Estimate conifer and hardwood covers separately.

Please note: These cover values should not include the coverage of regenerating tree species (i.e., tree seedlings and saplings).

**% Regenerating Tree:** The total foliar cover of seedlings and saplings, disregarding overlap of individual recruits. See seedling and sapling definitions below.

**%Shrub:** The total foliar cover (considering porosity) of all live shrub species disregarding overlap of individual shrubs.

**%Herbaceous:** The total cover (considering porosity) of all herbaceous species, disregarding overlap of individual herbs.

### **Height Class by Layer**

Modal height for conifer tree /hardwood tree, shrub, and herbaceous categories: Provide an estimate of height for each category listed. Record an average height value per each category by estimating the mean height for each group. Please use the following height intervals to record a height class: 01 = < 1/2m, 02=1/2-1m, 03 = 1-2 m, 04 = 2-5 m, 05 = 5-10 m, 06 = 10-15 m, 07 = 15-20 m, 08 = 20-35 m, 09 = 35-50 m, 10 => 50m.

### **Species List and Coverage**

- If mistletoe present add in what species it is living on
- Record absolute percent cover of dead tree species (can include saplings and seedlings)

**For rapid assessments,** list the 10-20 species that are dominant or that are characteristically consistent throughout the stand. These species may or may not be abundant, but they should be constant representatives in the survey. When different layers of vegetation occur in the stand, make sure to list species from each stratum. As a general guide, make sure to list at least 1-2 of the most abundant species per stratum.

**For relevés,** list all species present in the plot, using the second species list page if necessary.

**For both sample types,** provide the stratum:

T = Tree. A woody perennial plant that has a single trunk.

S = Shrub. A perennial, woody plant, that is multi-branched and doesn't die back to the ground every year.

H = Herb. An annual or perennial that dies down to ground level every year.

E = SEedling. A tree species clearly of a very young age that is < 1" dbh.

A = SApling. 1" - <6" dbh and young in age, OR small trees that are < 1" diameter at breast height, are clearly of appreciable age, and kept short by repeated browsing, burning, or other disturbance.

N = Non-vascular. Includes moss, lichen, liverworts, hornworts, cryptogamic crust, and algae.

Be consistent and don't break up a single species into two separate strata. The only time it would be appropriate to do so is when one or more tree species are regenerating, in which case the Seedling and/or Sapling strata should be recorded for that species. These may be noted on the same line, e.g.:

Strata	Species	%Cover	C
T/E/A	Quercus douglasii	40/<1/<1	

If a species collection is made, it should be indicated in the collection column with a "C" (for collected). If the species is later keyed out, cross out the species name or description and write the keyed species name in pen on the data sheet. Do not erase what was written in the field, because this information can be used if specimens get mixed up later. If the specimen is then thrown out, the "C" in the collection column should be crossed out. If the specimen is kept but is still not confidently identified, add a "U" to the "C" in the collection column (CU = collected and unconfirmed). In this case the unconfirmed species epithet should be put in parentheses [e.g. *Hordeum (murinum)*]. If the specimen is kept and is confidently identified, add a "C" to the existing "C" in the collection column (CC = Collected and confirmed).

Use Jepson Manual nomenclature. Write out the genus and species of the plant. Do not abbreviate. When uncertain of an identification (which you intend to confirm later) use parentheses to indicate what part of the determination needs to be confirmed. For example, you could write out *Brassica (nigra)* if you are sure it is a Brassica but you need further clarification on the specific epithet.

Provide the % absolute aerial cover for each species listed. When estimating, it is often helpful to think of coverage in terms of the following cover intervals at first:

<1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, >75%.

Keeping these classes in mind, then refine your estimate to a specific percentage. All species percent covers may total over 100% because of overlap.

Include the percent cover of snags (standing dead) of trees and shrubs. Note their species, if known, in the "Stand history, stand age and comments" section.

For rapid assessments, make sure that the major non-native species occurring in the stand also are listed in the space provided in the species list with their strata and % cover. For relevés, all non-native species should be included in the species list.

Also for relevés, you can record the <1% cover in two categories: r = trace (i.e., rare in plot, or solitary individuals) and + = <1% (few individuals at < 1% cover, but common in the plot).

**Unusual species:** List species that are locally or regionally rare, endangered, or atypical (e.g., range extension or range limit) within the stand. This field will be useful to the Program for obtaining data on regionally or locally significant populations of plants.

## INTERPRETATION OF STAND

**Field-assessed vegetation alliance name:** Name of alliance or habitat following the most recent CNPS classification system or the Manual of California Vegetation (Sawyer J.O., Keeler-Wolf T., and Evens, J. 2009). Please use scientific nomenclature, e.g., *Quercus agrifolia* forest. An alliance is based on the dominant or diagnostic species of the stand, and is usually of the uppermost and/or dominant height stratum. A dominant species covers the greatest area. A diagnostic species is consistently found in some vegetation types but not others.

**Please note:** The field-assessed alliance name may not exist in the present classification, in which case you can provide a new alliance name in this field. If this is the case, also make sure to state that it is not in the MCV under the explanation for “Confidence in alliance identification.”

**Field-assessed association name (optional):** Name of the species in the alliance and additional dominant/diagnostic species from any strata, as according to CNPS classification. In following naming conventions, species in differing strata are separated with a slash, and species in the uppermost stratum are listed first (e.g., *Quercus douglasii*/*Toxicodendron diversilobum*). Species in the same stratum are separated with a dash (e.g., *Quercus lobata*-*Quercus douglasii*).

**Please note:** The field-assessed association name may not exist in the present classification, in which you can provide a new association name in this field.

**Adjacent Alliances/direction:** Identify other vegetation types that are directly adjacent to the stand being assessed by noting the dominant species (or known type). Also note the distance away in meters from the GPS waypoint and the direction in degrees aspect that the adjacent alliance is found

(e.g., *Amsinckia tessellata* / 50m, 360°N *Eriogonum fasciculatum* /100m, 110°).

**Confidence in Identification:** (L, M, H) With respect to the “field-assessed alliance name”, note whether you have L (=Low), M (=Moderate), or H (=High) confidence in the interpretation of this alliance name.

**Explain:** Please elaborate if your “Confidence in Identification” is low or moderate. Low confidence can occur from such things as a poor view of the stand, an unusual mix of species that does not meet the criteria of any described alliance, or a low confidence in your ability to identify species that are significant members of the stand.

**Phenology:** Indicate early (E), peak (P) or late (L) phenology for each of the strata.

**Other identification problems or mapping issues:** Discuss any further problems with the identification of the assessment or issues that may be of interest to mappers. Note if this sample represents a type that is likely too small to map. If it does, how much of the likely mapping unit would be comprised of this type. For example: “this sample represents the top of kangaroo rat precincts in this general area, which are surrounded by vegetation represented by CARR000x; this type makes up 10% of the mapping unit.” Depending on who mapped polygon (Calveg, etc); we should denote that information here.

**Is polygon >1 type:** Yes / No (circle one): In areas that have been delineated as polygons on aerial photographs/imagery for a vegetation-mapping project, assess if the polygon is mapped as a single stand. “Yes” is noted when the polygon delineated contains the field-assessed alliance and other vegetation type(s), as based on species composition and structure. “No” is noted when the polygon is primarily representative of the field-assessed alliance.

**If yes, explain:** If “Yes” above, explain the other vegetation alliances that are included within the polygon, and explain the amount and location that they cover in the polygon.



**Other CNDDDB/Whitebark Pine (WBP) monitoring Data:**

Trees/stems are assessed within a representative portion of the stand (using a specific radius or area for averaging).

**Mountain Pine Beetle (MPB) Level: Should equal 100%.**

Note the level of mountain pine beetle attack using the following:

- 0 = No evidence of attack or beetle pitch tubes or unknown
- 1 = less than 5 observable beetle pitch tubes ('hits')
- 2 = less than 50% of the bole is attacked; sporadic pitch tubes spread on most parts of the bole or several localized areas with a high density (>10) pitch tubes
- 3 = greater than 50% of the bole is attacked; numerous pitch tubes spread on many parts of the bole

**% of WBP Cones (female only): Should equal 100%.**

Record the number of cones in the tree/stem using the following numeric system:

- 0 = no cones
- 1 = 1 to 10 cones
- 2 = 11 to 100 cones
- 3 = greater than 100 cones

**Total # WBP individuals or clumps and size (CNDDDB):**

The number of individuals observed/detected during assessment. This should be recorded as clumps (or # of stems within # of clumps) per defined area (square meters, hectares, acres, etc.).

**Phenology of WBP (CNDDDB): Should equal 100%.**

The average percent of WBP that is vegetative, flowering (nascent female cones) and/or fruiting (mature female cones).

**% WBP mortality:**

These percentages are for mortality of trees/stems from mountain pine beetle (MPB) or white pine blister rust (WPBR); 'Other' can be % mortality from both MPB and WPBR; including WPBR mortality on other species E.g. WPBR-PIMO/PIBA 5% (white pine blister rust on *Pinus monticola* or *Pinus balfouriana* at 5% cover) or unknown causes.

**Overall site/occurrence quality/viability (site + population) (CNDDDB):**

Is the likely persistence of the occurrence into the future Excellent, Good, Fair, or Poor? This is an assessment of the overall viability of this occurrence. Both the quality & condition of the site and of the occurrence must be considered when scoring. Take into account population size, demography, viability over time, site condition, and any disturbances. And also see additional characteristics at: <http://www.natureserve.org/explorer/eorankguide.htm>

**Determination of WBP:** Please indicate how the species identification was determined.

**CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)**

Relevé or Rapid Assessment (circle one) (Revised June 28, 2013)

For Office Use:	Final database #:	Final vegetation type name:	Alliance Association
-----------------	-------------------	-----------------------------	----------------------

**I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION**

Polygon/Stand #:	Air photo:	Date:	Name(s) of surveyors (circle recorder):
------------------	------------	-------	---

GPS wypt #: \_\_\_\_\_ GPS name: \_\_\_\_\_ Datum: \_\_\_\_\_ or NAD83. Bearing, left axis at SW pt \_\_\_\_\_ (degrees) of Long / Short side  
 UTM E \_\_\_\_\_ UTM N \_\_\_\_\_ Zone: 10 / 11 (circle one) Error: ± \_\_\_\_\_ ft / m / pdop  
 GPS within stand? Yes / No If No, cite from waypoint to stand, distance \_\_\_\_\_ (meters) & bearing \_\_\_\_\_ (degrees)

Elevation: \_\_\_\_\_ ft / m Camera Name/Photograph #'s: \_\_\_\_\_

Stand Size (ac/ha): <1, 1-5, >5 ac | \_\_\_\_\_ ha Plot Size (m<sup>2</sup>): 10 / 100 / 400 | Plot Shape \_\_\_\_\_ x \_\_\_\_\_ m or Circle Radius \_\_\_\_\_ m  
 Exposure, Actual °: \_\_\_\_\_ NE NW SE SW Flat Variable All | Steepness, Actual °: \_\_\_\_\_ 0° 1-5° 5-25° > 25

Topography: Macro: top upper mid lower bottom | Micro: convex flat concave undulating  
 Geology code: \_\_\_\_\_ Soil Texture code: \_\_\_\_\_ | Upland or Wetland/Riparian (circle one)

% Surface cover: (Incl. outcrops) (>60cm diam) (25-60cm) (7.5-25cm) (2mm-7.5cm) (Incl sand, mud)  
 H20: \_\_\_\_\_ BA Stems: \_\_\_\_\_ Litter: \_\_\_\_\_ Bedrock: \_\_\_\_\_ Boulder: \_\_\_\_\_ Stone: \_\_\_\_\_ Cobble: \_\_\_\_\_ Gravel: \_\_\_\_\_ Fines: \_\_\_\_\_ =100%

% Current year bioturbation \_\_\_\_\_ Past bioturbation present? Yes / No | Fire evidence: Yes / No (if yes, explain below)  
 Habitat description, surrounding land use, comments (CNDDDB):

\_\_\_\_\_

\_\_\_\_\_

Disturbance / Intensity (L,M,H) \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_ WBP Impact 39 / \_\_\_\_\_ / \_\_\_\_\_ 40 / \_\_\_\_\_ / \_\_\_\_\_

**II. HABITAT AND VEGETATION DESCRIPTION**

Tree DBH: **T1** (<1" dbh), **T2** (1-6" dbh), **T3** (6-11" dbh), **T4** (11-24" dbh), **T5** (>24" dbh), **T6** multi-layered (T3 or T4 layer under T5, >60% cover)  
 Shrub: **S1** seedling (<3 yr. old), **S2** young (<1% dead), **S3** mature (1-25% dead), **S4** decadent (>25% dead)  
 Herbaceous: **H1** (<12" plant ht.), **H2** (>12" ht.) % NonVasc cover: \_\_\_\_\_ % Vasc Veg cover: \_\_\_\_\_  
 % Cover - Conifer tree / Hardwood tree: \_\_\_\_\_ / \_\_\_\_\_ Regenerating Tree: \_\_\_\_\_ Shrub: \_\_\_\_\_ Herbaceous: \_\_\_\_\_  
 Height Class - Conifer tree / Hardwood tree: \_\_\_\_\_ / \_\_\_\_\_ Regenerating Tree: \_\_\_\_\_ Shrub: \_\_\_\_\_ Herbaceous: \_\_\_\_\_  
 Height classes: 01=<1/2m 02=1/2-1m 03=1-2m 04=2-5m 05=5-10m 06=10-15m 07=15-20m 08=20-35m 09=35-50m 10=>50m

Species, Stratum, and % cover. Stratum categories: T=Tree, S = Shrub, H= Herb, E = SEedling, A = SApling, N= Non-vascular.  
 % cover intervals for reference: <1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, 75%.

Strata	Species	% dead	% cover	C	Strata	Species	% dead	% cover	C

Other rare taxa in stand (CNDDDB) \_\_\_\_\_

**III. INTERPRETATION OF STAND**

Field-assessed vegetation alliance name: \_\_\_\_\_

Field-assessed association name (optional): \_\_\_\_\_

Adjacent alliances/direction: \_\_\_\_\_ / \_\_\_\_\_, \_\_\_\_\_ / \_\_\_\_\_

Confidence in alliance identification: L M H Explain: \_\_\_\_\_

Other identification or mapping information: \_\_\_\_\_ Phenology (E,P,L): Herb \_\_\_\_\_ Shrub \_\_\_\_\_ Tree \_\_\_\_\_

Is poly >1 type: Yes / No If yes, explain: \_\_\_\_\_

**CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)  
Other CNDDDB/Whitebark Pine Monitoring Data:**

Polygon/Stand #:
MBP Level: 0=no attack _____ % 1=>5 hits _____ % 2=<50% of bole attacked _____ % 3=>50% of bole attacked _____ %
Avg % of WBP Cones: No cones _____ % 1-10 cones _____ % 11-100 _____ % >100 _____ %
Total # individuals or clumps (WBP) and size (CNDDDB) _____ # per _____ hectares
Phenology of WBP (CNDDDB): Vegetative _____ % Flowering (cones) _____ % Fruiting (cones) _____ %
%WBP mortality: MPB _____ % WPBR _____ % Other: _____ % _____ %
Overall site/occurrence quality/viability (site + population) (CNDDDB): <input type="checkbox"/> Excellent <input type="checkbox"/> Good <input type="checkbox"/> Fair <input type="checkbox"/> Poor
Determination of WBP: Keyed _____ By another person (name) _____ Compared with photo/drawing _____ Other _____

Sample Rapid Assessment from the Boulder Peak Region - Marble Mountain Wilderness

**CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)  
Other CNDDDB/Whitebark Pine Monitoring Data:**

Polygon/Stand #: BP01
MBP Level: 0=no attack <u>70</u> % 1=>5 hits <u>—</u> % 2=<50% of bole attacked <u>—</u> % 3=>50% of bole attacked <u>30</u> %
Avg % of WBP Cones: No cones <u>70</u> % 1-10 cones <u>20</u> % 11-100 <u>10</u> % >100 <u>—</u> %
Total # individual clumps (WBP) and size (CNDDDB) <u>15</u> # per <u>2</u> hectares
Phenology of WBP (CNDDDB): vegetative <u>70</u> % flowering (cones) <u>20</u> % fruiting (cones) <u>20</u> %
%WBP mortality: MPB <u>30</u> % WPB <u>0</u> % Other: <u>—</u> % _____ %
Overall site/occurrence quality/viability (site + population) (CNDDDB): <input type="checkbox"/> Excellent <input type="checkbox"/> Good <input checked="" type="checkbox"/> Fair <input type="checkbox"/> Poor
Determination of WBP: Keyed _____ By another person (name) _____ Compared with photo/drawing _____ Other _____

White fir / Shasta fir ATTACK by fir engraver with 20% mortality

WWP (P. monticola) 1 dead MPB.

CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)  
 Relevé or Rapid Assessment (circle one) (Revised June 28, 2013)

For Office Use: Final database #: Final vegetation type name: Alliance Association

I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION

Polygon/Stand #: BPO1 Air photo: Date: 9/20/13 Name(s) of surveyors (circle recorder): KARKMAN

GPS wypt #: GPS name: Datum: or NAD83. Bearing, left axis at SW pt (degrees) of Long / Short side  
 UTME 492814 UTMN 4604449 Zone: 10/11 (circle one) Error: ±2.3 ft/m / pdop  
 GPS within stand? Yes/No If No, cite from waypoint to stand, distance (meters) & bearing (degrees)

Elevation: 1985 ft/m Camera Name/Photograph #'s:

Stand Size (ac/ha): <1, 1-5, >5 ac/ ha Plot Size (mz): 10 / 100 / 500 Plot Shape x m or Circle Radius 12 m  
 Exposure, Actual °: NE NW SE SW Flat Variable All | Steepness, Actual °: 3° 0° 1-5° 5-25° >25

Topography: Macro: top upper mid lower bottom | Micro: convex flat concave undulating  
 Geology code: GAbbr Soil Texture code: Upland or Wetland/Riparian (circle one)

% Surface cover: (Incl. overcrops) (>60cm diam) (25-60cm) (7.5-25cm) (2mm-7.5cm) (Incl sand, mud)  
 H20: 50 BA Stems: 30 Litter: 57 Bedrock: Boulder: 10 Stone: 5 Cobble: 5 Gravel: 5 Fines: 10 =100%

% Current year bioturbation Past bioturbation present? Yes/No | Fire evidence: Yes No (if yes, explain below)  
 Habitat description, surrounding land use, comments (CNDDDB):

On edge of creek, 1/2 mile below lower Wright Lake (on trail)  
 meadow edge - RIPARIAN. Fire evidence from 20+ years ago  
 now with white fir encroaching.

Disturbance / Intensity (L,M,H) Grazing / L WBP Impact 39 1M, 50% 40 16, 30%

II. HABITAT AND VEGETATION DESCRIPTION

Tree DBH: T1 (<1" dbh), T2 (1-6" dbh), T3 (6-11" dbh), T4 (11-24" dbh), T5 (>24" dbh), T6 multi-layered (T3 or T4 layer under T5, >60% cover)  
 Shrub: S1 seedling (<3 yr. old), S2 young (<1% dead), S3 mature (1-25% dead), S4 decadent (>25% dead)  
 Herbaceous: H1 (<12" plant ht.), H2 (>12" ht.) % NonVasc cover: 35 % Vasc Veg cover: 65  
 % Cover - Conifer tree / Hardwood tree: 15 / Regenerating Tree: 5 Shrub: 25 Herbaceous: 20  
 Height Class - Conifer tree / Hardwood tree: 05 / Regenerating Tree: 02 Shrub: 02 Herbaceous: 01  
 Height classes: 01=<1/2m 02=1/2-1m 03=1-2m 04=2-5m 05=5-10m 06=10-15m 07=15-20m 08=20-35m 09=35-50m 10=>50m

Species, Stratum, and % cover. Stratum categories: T=Tree, S=Shrub, H=Herb, E=SEedling, A=SApling, N=Non-vascular.  
 % cover intervals for reference: <1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, 75%

Strata	Species	% dead	% cover	C	Strata	Species	% dead	% cover	C
T	Whitebark	30%	5%		S	Servicberry		3%	
T	White fir	10	10%		S	Ribes sp.		4%	
T	Shasta fir		3%		H	Grass sp.		10%	
T	Western white pine (1)	dead-1 year			H	Worm sp		5%	
S	SALIX sp.		5%		H	elderberry		5%	
S	Snowberry		5%		T	FOXTAIL pine - 4 new		3%	
S	VACINIUM sp.		5%			reg trees			
S	Sorbus sp.		3%						

Other rare taxa in stand (CNDDDB)

III. INTERPRETATION OF STAND

Field-assessed vegetation alliance name: White fir - WBP  
 Field-assessed association name (optional): meadow to  
 Adjacent alliances/direction: EAST  
 Confidence in alliance identification: L (M) H Explain:

Other identification or mapping information: Along trail Phenology (E,P,L): Herb Shrub Tree

Is poly >1 type: Yes / No If yes, explain:

# Appendix 2: Recommended Protocols for Future Work

## Whitebark Pine Inventory and Monitoring Plot Protocol

Revised: May 16, 2013

### Introduction:

This protocol was developed collaboratively by the USFS Region 5 Ecology Program and Forest Health Protection Program to provide inventory and status-and-trend monitoring data in stands dominated by whitebark pine (*Pinus albicaulis*) or lodgepole pine (*P. contorta*) with whitebark pine as a codominant species. It also focuses on stands that have experienced recent tree mortality related to insects (mountain pine beetle) or diseases (white pine blister rust). This protocol was developed using elements of the Regional Ecology Program post-fire regeneration monitoring protocol and the Forest Health Protection Whitebark Pine Monitoring Plot Protocol for the Warner Mountains, Modoc National Forest (the FHP protocol is based on *Greater Yellowstone Whitebark Pine Monitoring Working Group [GYWPMWG] Interagency Whitebark Pine Monitoring Protocol for the Greater Yellowstone Ecosystem 2007*).

### I. Site attributes:

- Record the location (geographic or watershed scale), site (topographic scale), and plot number (micro-scale).
- Use a GPS device to locate plot center - take care to avoid biasing the location.
- Monument permanent plots (established for monitoring rather than inventory) with 2-foot long rebar driven approximately 1.5 ft into the ground at plot center. Label with plot number and mount safety cap. Include brief notes of plot location using distinctive landscape features, if any.
- Record the date that data were collected.
- Record the crew names of the people collecting the data.
- Establish a plot with a radius of 12.6 m, which is approximately 0.05 ha (0.124 acres). Flag four places around the perimeter for reference.
- Record the dominant tree species present.
- Take one photograph from a point 12.6 m south of the plot center, looking north. Make sure you have something (pin flag) at plot center so it can be relocated. using the photo. Take another photograph from a point 12.6 m north of the plot center, looking south (toward pin flag). Record both photo numbers.
- Record the average slope of the plot in percent (use clinometer).
- Record the average aspect of the plot in degrees (use compass [make sure you have the right declination])

### II. Vegetation and ground cover attributes

- Estimate the cover (%) of: *basal vegetation* (i.e. the area covered by the bases of tree boles, shrub stems, herbs), *litter*, *bare ground*, *rock* (>2 mm diameter), and *woody debris* (>3 inches [7.5 cm] diameter), summing to 100% (imagine chopping off all vegetation at ground level, what do you have left?; it is rare for basal vegetation to be more than 5%, unless there are trees or many large shrubs in the plot). Record cover values to nearest 5%, using 0.5% as trace cover
- Estimate vegetation cover to nearest 1% (1-10% cover), 5% (10-30%) or 10% (30-100%):

- % Overstory veg cover = cover of plants >2 m in height (trees and tall shrubs; this is a snapshot of total canopy cover taken from above, i.e. it is the % of the plot that has trees/tall shrubs covering it. Tree/tall shrubs growing completely beneath other trees/tall shrubs are not counted as they cannot be seen from above) (see Figure 1)
  - Estimate cover of live trees and tall shrubs
  - Also estimate % dead cover (trace circles around the remnants of dead trees >2 m in height). When this value is added to the live cover it should give us an estimate of the total pre-beetle mortality live cover.
- % Shrub and Herbaceous plant cover = cover of understory vegetation <2 m in height (this is a snapshot of total understory cover taken from above, i.e. it is the % of the plot that has understory vegetation covering it. Understory plants growing completely beneath other plants are not counted, as they cannot be seen from above) (see Figure 1).
- Record separately the cover of aspen <3 m in height. Aspen >3 m height should be recorded as overstory cover.




Fig. 1. Circular plot, with four species of understory plants (colored). The blue crosshairs are added to aid in estimating cover. The understory vegetation cover is about 64% (the total plot area minus the area that is not covered by live vegetation). The gray species (shrub) has 49% cover, the dark blue species (shrub) has 17% cover, the light blue species (grass) has 4%, and the orange species (forb) has 6% (each orange circle is 1% in this case). Due to plant overlap, summing the different species' cover values gives a value that is larger than the total understory vegetation cover (76% vs. 64%). Overall shrub cover in this plot is 58% (two shrub species, subtracting overlap; summed up [i.e., ignoring overlap], the two species have 66% cover between them). Herb cover is 9.5%; forb cover is 6%; grass cover is 4%. Each shrub species will have its own cover entered in the species-cover section of the datasheet. Cover is measured by drawing a line around the outside of the plant canopy, ignoring gaps that may be found within the perimeter. For plots of this size (500 m<sup>2</sup>), your clipboard is about 0.015% of the plot area.

### III. Basal area, snags, and litter depth

- Use the basal area gauge (20 factor) to record the basal area of live and dead whitebark pine and other tree species (e.g., lodgepole pine) in the stand
  - Swinging the gauge around the plot center, tally the number of trees that are larger than the 20 factor aperture. Count live and dead trees separately for each species.
- Record the species and dbh of any older snags (>7 years; **prior to 2005**) in the plot. Only record snags that are >1.37 m tall.
- Measure litter depth at 3 locations midpoint between plot center and plot perimeter in 3 directions (0°, 120°, and 240°).

### IV. Tree regeneration attributes

- Tally the number of seedlings and saplings (trees less than 7.6 cm dbh of each tree species for each age class)
  - Use a separate row for each species and basal cluster (see below).
  - Count the number of live and dead stems arising from each seedling or sapling cluster. Clusters are defined using the following two criteria:
    - Stems are less than 10 cm at the base from the cluster of other stems
    - Diameter of stem (saplings only) must not exceed 25% of next largest stem in the cluster
  - Determine minimum age by counting the bud scars, subtracting the current year
  - Record dbh for saplings (>1.37 m height) only
  - In the Health Code column, note the number of seedlings or saplings in each cluster that exhibit health issues and include the appropriate health codes for these numbers (e.g., 2-a). Health codes include:
    - C = cankers or stem swelling
    - SC = stalactiform canker (*P. contorta* only) as spindle-like in middle of bole
    - P = pitching
    - F = branch flagging
    - S = needle spots
    - T = twig beetle sign (e.g. terminal branch flagging and pitch tubes)
    - 2 = secondary beetle
    - M = dwarf mistletoe
    - R = native rusts – take photo and collect sample when available
    - H = sapsucker/woodpecker holes
    - A = aecia (i.e., rust fruiting body) or aeciospores
      - Take closeup photos of any branches displaying aecia and consider collecting samples for laboratory identification
      - Aecia could be a sign on WPBR or a native rust
  - Record the height for the tallest individual seedling of each species

## V. Understory vegetation attributes

- Measure the modal height and overall cover for the four most common (by % cover) shrub and herbaceous plant species in the plot. Additional understory species may be noted in the Notes section. Especially note the presence of species in the genera *Ribes*, *Castilleja*, and *Pedicularis* (WPBR secondary hosts)
  - Measure cover to nearest 5%, 0.5% = trace cover
  - Modal height is the most common height

## VI. Notes section

Items of interest to record in the notes section:

- If fire scars or other evidence of fire are in the plot
- If plot is located on a unique (non-granitic) substrate (e.g., pumice soils)
- If plot has been treated in some way - specify
- If non-native species are on plot or adjacent to plot - specify
- If other mortality agents (insects, diseases) are present - specify
- If WPBR, mountain pine beetle, twig beetle, or other potential mortality agents are observed adjacent to plot but not recorded within the plot
- If conifer stumps are present from trees that may have parented seedlings before they were cut
- Additional understory species if more than four shrubs or herbaceous plants
- Other notes?

## VII. Tree attributes

For all trees (>7.6 cm dbh) record the following information

- Species ID, and number live and dead stems in each cluster. Tree clusters are defined by stems that are less than 1 m at the base from the cluster of other stems of similar size (diameter and height).
- Individual stems growing in close proximity will be defined as individual tree stems or branches using the following GYWPMWG (2007) criteria:
  - 1) There must be a discernible growth groove that separates that stem from other stems of the tree.
  - 2) The diameter of a given stem must be more than 25% of the diameter of the largest stem.
  - 3) The stem must be less than one foot from the “mother” tree to which it is associated. Otherwise it is to be considered as a separate seedling, sapling, or tree.
  - 4) The angle of the stem in question must be no less than a 45° angle from the main stem.
- For each cluster, provide a consecutively-numbered cluster ID number. For each stem within a cluster, provide a stem ID value.
- Record the dbh of all live and dead stems in each cluster.



- For monumented monitoring plots, nail aluminum tags to all live trees that are counted with the basal area gauge (20 factor) sweep. Begin consecutively-numbered tags at the northernmost tree proceeding clockwise. Nail tags at dbh so that each tag faces plot center, leaving approximately one inch of space between the nail head and the tree bole in order to avoid damage during tree growth. Enter tag numbers in the Notes column.
- Note with a checkmark whether live basal sprouts are present for a given tree cluster. These basal sprouts are defined as smaller-diameter (typically <7.6 cm dbh) live stems located at the base and often surrounding clusters of larger live and dead tree clusters (typically with stems exceeding 20 cm dbh).
- Note the % of live crown in the stem (largest live is the default) in increments of 10% using the following coding system: 1 = 10%, 2 = 20%, 3 = 30%, etc.
- Note the level of mountain pine beetle attack using the following:
  - 0 = No evidence of attack or beetle pitch tubes or unknown
  - 1 = less than 5 observable beetle pitch tubes ('hits')
  - 2 = less than 50% of the bole is attacked; sporadic pitch tubes spread on most parts of the bole or several localized areas with a high density (>10) pitch tubes
  - 3 = greater than 50% of the bole is attacked; numerous pitch tubes spread on many parts of the bole
- Estimate the time since mountain pine beetle attack based on the following system:
  - 0 = less than one year since attack (occurred during current season); typically little sign of crown discoloration or dead needles but evidence of beetle attack
  - 1 = approximately one year since attack (last season); crown shows significant density of dead or dying needles (substantial portion of crown contains brown/orange colored needles)
  - 2 = two years since attack; entire crown consists of dead (brown/orange colored) needles that are mostly intact
  - 3 = three years since attack; most but not all of dead needles have fallen from crown, with few clusters of dead needles retained
  - 4 = four to seven years since attack; no dead needles retained in crown; smaller branches may have broken off and fallen, with most larger to medium branches retained
- Record the number of cones in the tree using the following numeric system:
  - 0 = no cones
  - 1 = 1 to 10 cones
  - 2 = 11 to 100 cones
  - 3 = greater than 100 cones
- Record the health code for each tree using the codes listed under the tree regeneration attributes section (see above)
- In Notes column record any remarkable observations pertaining to a tree or tree cluster, including:
  - lightning or fire scars

- evidence of other damage caused by wildlife, humans, or unknown causes
- Other notable features or observations

#### VIII. Seed-Caching Wildlife Point Counts (optional, if time permits)

- At end of vegetation sampling period, return to each plot and spend 5 minutes noting any visual or auditory sign of Clark's Nutcracker (*Nucifraga columbiana*), Douglas' squirrel (*Tamiasciurus douglasii*), lodgepole chipmunk (*Neotamias speciosus*), golden-mantled ground squirrel (*Callospermophilus lateralis*), or any other seed-eating species within 50 m of each side of transect. Record plot number, observer, time and date of survey, and number of each species observed at each sample point (i.e., plot).
- Note any observations of seed caching, seed dispersal, or seed predation during survey period.
- Record data on separate field notebook

# 2018 Update

## Whitebark Pine Pilot Fieldwork Report

### Klamath National Forest



By Michael Kauffmann<sup>1</sup>, Raphaela Floreani Buzbee, Jennifer Buck-Diaz,  
Julie Evens, and Sydney Magner

In collaboration with:

Erin Lonergan, Forest Botanist, Modoc National Forest

Diane Ikeda, Regional Botanist, Pacific Southwest Region, USDA Forest Service

**NOVEMBER 2018**

---

1. Kauffmann, Michael E., PO Box 777, Bayside, CA 95524, [michaielekauffmann@gmail.com](mailto:michaielekauffmann@gmail.com)



CALIFORNIA  
NATIVE PLANT SOCIETY



ABOVE: *Whitebark pine on the summit of China Mountain.*

## Methods and Materials

The California Native Plant Society (CNPS) obtained updated GIS data from various sources including the USFS Pacific Southwest - Region Remote Sensing Lab's CALVEG maps (USFS 2018), and California Department of Fish and Wildlife (CDFW) - High Elevation Species Sightings Sierra Nevada (CDFW 2018). In addition, we used the updated species profile from the CNPS Rare Plant Program, which maps the distribution of whitebark pine using many sources such as herbarium specimens and other observational data (CNPS 2018).

CNPS reviewed the CNPS/CDFW Vegetation Rapid Assessment protocol used in the 2013 pilot project for evaluating whitebark pine vegetation and insect/disease impacts. We made minor edits and clarification to the data sheet and protocol in order to collect standardized data across the five forests. A reconnaissance (recon) form was used to gather simplified, general information about stands of vegetation, especially for WBP stands that were near or adjacent to stands already sampled by a full Rapid Assessment.

## Notes on Presence and Alliance

A vegetation alliance is a category of vegetation classification which describes repeating patterns of plants across a landscape. Each alliance is defined by plant species composition, and reflects the effects of local climate, soil, water, disturbance, and other environmental factors. Alliances are commonly used in vegetation mapping.

Presence indicates that whitebark pine can be found but is not the dominant vegetation pattern.



*An example of a whitebark pine alliance, just south of the Boulder Peak summit.*

The following areas, prioritized for future work after our 2013 surveys, were visited in the summer of 2018.

- Scott River District



1. Assess the population on China Mountain
2. Ground truth and create a better map for the Russian Wilderness area.
3. Ground truth the Heather Lake Pluton, between Man Eaten and Cliff lakes and create a better map for the Boulder Peak Region.
4. Survey Gooseneck and Willow Creek Mountains.

### 1. China Mountain Region

This area offers the only opportunity to drive to a whitebark pine in the Klamath Mountains. That being said, most of the trees are on the extensive ridgeline radiating in all directions from China Mountain. Five polygons show the whitebark pine alliance to be stronger than previously mapped, making this an important holdout for the species regionally.

Trees in this region, similar to those on the Boulder Peak Ridgeline, are showing the highest current impacts of mountain pine beetle infestation. The three surveys in the PIAL Alliance showed between 15-40% mortality from beetle kill.

California endemic foxtail pines (*Pinus balfouriana*) are also found mixed with whitebark here. Because this species is also a 5-needle pine, it is affected by the same pathogens as whitebark. Thus, trees here show similar mortality, but at a slightly lower rates—possibly due to the rarity of this species and beetles preference, or familiarity, for whitebark pine.




## 2. Russian Wilderness

This region offers a narrow ridgeline with an average elevation high enough to support whitebark pine. This ridgeline, between 7,500-8,000 feet, is the divide between the Salmon and Scott rivers in the heart of the Russian Wilderness. We accessed this area by the Bingham Lake trailhead.

Evidence of both blister rust and bark beetles was found. In fact, a survey captured one of only a handful of instances of mortality caused by blister rust.

To the west of the Pacific Crest Trail, near Siphon Lake, there are a few isolated occurrences of individual trees, or clumps of fewer than 10 trees, that appear to be more recently established by nutcracker dispersal.

Future work here should include mapping the ridges above Lake Albert (circled in pink on the map to the right). The layers for the southern wilderness are in good shape.





### 3. Northern Marble Mountains

Because of issues with smoke and fire in other areas of the forest, I pursued more ground-truthing in the Boulder Peak Region which proved informative when coupled with the 2013 surveys. While this extensive series of whitebark pine groves is one of the largest (probably the largest) in the Klamath Mountains along the ridgelines surrounding Boulder Peak, specimens also grow at some of the lowest elevations in the range (~6,500 feet). The species did not extend into as many drainages as I had previously predicted around Boulder Peak.

The spine of this extensive high elevation ridgeline was typically a whitebark pine or whitebark pine-foxtail pine alliance. These trees grew along the relatively flat ridgeline or on the south-southeast-southwest slopes. In these alliance stands mortality from mountain pine beetles is approaching 30% in some areas with small occurrences of active kills from this year and recent previous years. There is also evidence of white pine blister rust across this entire region.

To the south of Boulder Peak, I visited a few of the high points near Man Eaten Lake. While I did find WBP, they only occurred at the summit of one high point (see circle on map). This stand is experiencing 70% mortality from both MPB and a fire that burned up the ridge from Wooley Creek in 2017. There is a possibility that more PIAL could be found to the south around Wooley Lake, though my field notes from 2008 make no mention of the species there.



## Shasta North

### • Goosenest Ranger District

#### 1. Ground-truth Willow Creek Mountain and Goosenest Peak

##### **Goosenest**

This volcano summit is quite cool. The well established trail starts in the middle of nowhere, immediately in the presences of whitebark pine. Within the Shasta fir alliance, trees show low levels of mortality (<5%) from occasional mountain pine beetle kill. Near and within the summit crater, whitebark pine alliance is the norm, except on the north slopes within the crater where mountain hemlock is the alliance.

Within the crater, almost every tree shows signs of needle browning. I hypothesize this is due to aridification of the summit due to decreased snowpack and increased annual average temperature. This would be an interesting location for a long term monitoring plot.

##### **Willow Creek**

This volcano summit, close to the Little Shasta Meadow Botanical Area, is a quick climb from the Willow Creek Mountain Road. The summit revealed several small stands of whitebark pine alliance. The tree was present mixed with Shasta fir and mountain hemlock. Both blister rust and mountain pine beetles were present, but at low levels (<5%). This stand was previously undocumented.






Table 1: Area by forest as mapped by Michael Kauffmann.

Forest		2014 Presences (acres)	2018 Presences (acres)	2018 Alliance (acres)
Klamath National Forest	total	<del>9,198</del>	5,708.6	659.5
Shasta-Trinity National Forest	total	22,039	Did not map for alliance in 2013, only presence.  Not revised in 2018.	
Lassen National Forest (including Lassen N.P.)	total	12,095		
Modoc National Forest	total	<del>26,858</del>	23,472.6	1,357.5
Total acreage in the four forests of Northern California		<del>70,906</del>	63,315	--

Notes on the area changes:

Major revisions were made to the polygons in the northeastern Marble Mountain Wilderness. Based on initial surveys, particularly on the north slopes above the Scott River, I expected the species to be present at lower elevations. While this was occasionally the case, the pattern did not hold true across the entirety of the Boulder Peak ridgeline and the presence of whitebark pine was much smaller in area than predicted. The area in the Marble Mountains near Man Eaten Lake is also quite high in average elevation. I visited this area in 2004 and found a few trees along a lower elevation ridgeline. This led me to predict more in the area, however this did not hold true. In fact, the two areas I did find the trees were less than 5 acres in size. In all, the variability of terrain, steep and narrow ridgelines, as well as large boulder fields limits the habitat for whitebark pine in this area of the Marble Mountains. Other decreases in predicted area were found in the southern Russian Wilderness for the same reasons.

## Goosenest Ranger District



*Whitebark pines in the summit crater of Goosenest Mountain. Notice slight needle browning in all trees.*



*Whitebark pines on the south-facing summit of Goosenest Mountain.*



*Whitebark pine alliance on the summit of Willow Creek Mountain.*


## Scott River District



*Whitebark pines mortality from mountain pine beetles on the ridge north of China Mountain.*



*Sparse stands of PIAL in the southern Russian Wilderness are affected by both MPB and WPBR.*



*Near Man Eaten lake in the Marble Mountains, a small stand of PIAL was burned in 2017 - now with 70% mortality.*

## Recommendations for future KNF work:

### • Gooseneck Ranger District

1. Survey Garner Mountain
2. Look closer at the West Haight Mountain stands (particularly along the east ridge) and possibly pursue the designation of a botanical area.

### • Scott River

1. Survey and map the “Big Ridge” between Black Marble Mountain and King’s Castle in the Marble Mountain Wilderness.
2. Set up long term monitoring plots in the Boulder Peak Region. This is one of the most extensive stands of whitebark pine in the Klamath Mountains and most likely serves as a “feeder” population for the smaller mountain-top stands nearby. This area is critical to the future of whitebark in the Klamath Mountains.
3. Ground truth the higher peaks around Upper Albert and Big Blue lakes. I’m actually not 100% the species is here, but they should be!
4. Set up a permanent plot on South China Mountain.
5. Work with Shasta-Trinity to ground-truth the Cory Peak Botanical and Geological area to verify or nullify species occurrence.

### • Salmon River Ranger District

1. Work with the Weaverville and Big Bar RDs on the Shasta-Trinity to map and ground-truth the extent of WBP along the Stuarts Fork-Salmon Divide (this is some steep country!).
2. I believe that WBP could occur in the Dorleska Mine region in the Big Flat Region. This area should be ground-truthed and mapped

### • Across the Klamath National Forest

1. Create a map to target areas where encroachment from firs and hemlocks is an issue and consider managing for this problem



*China Mountain as seen from the south-facing slopes above Crater Lake.*


Table 2. Rapid Assessment summary, Klamath National Forest

DbaseID	County	Ranger District	Wilderness	Site Name	Alliance	Estimated Pct Cover PIAL	Altitude (m)	Impacts
WBP0900	Siskiyou	Scott River	n/a	China Mountain	<i>Pinus balfouriana</i>	10	2484	MPB (15%)
WBP0901	Siskiyou	Mt. Shasta	n/a	China Mountain	<i>Pinus balfouriana</i>	7	2497	MPB (20%)
WBP0902	Siskiyou	Scott River	n/a	China Mountain	<i>Pinus albicaulis</i>	10	2466	MPB (40%)
WBP0903	Siskiyou	Scott River	n/a	China Mountain	<i>Abies magnifica</i>	4	2382	None
WBP0904	Siskiyou	Goosenest	n/a	Goosenest Mountain	<i>Pinus albicaulis</i>	3	2438	None
WBP0905	Siskiyou	Goosenest	n/a	Goosenest Mountain (Private)	<i>Pinus albicaulis</i>	9	2491	MPB (15%)
WBP0906	Siskiyou	Goosenest	n/a	Goosenest Mountain	<i>Tsuga mertensiana</i>	5	2063	None
WBP0907	Siskiyou	Goosenest	n/a	Goosenest Mountain	<i>Tsuga mertensiana</i>	5	2295	MPB (20%)
WBP0908	Siskiyou	Goosenest	n/a	Willow Creek Mountain	<i>Abies magnifica</i>	6	2371	MPB (5%), WPBR (5%)
WBP0909	Siskiyou	Goosenest	n/a	Willow Creek Mountain	<i>Pinus albicaulis</i>	7	2385	None
WBP0921	Siskiyou	Scott River	n/a	Grizzly Peak	<i>Tsuga mertensiana</i>	4	2311	WPBR (5%)
WBP0922	Siskiyou	Salmon River	Russian	Trail Creek	<i>Tsuga mertensiana</i>	2	2233	None
WBP0923	Siskiyou	Scott River	Marble Mountain	Marble Mountains	<i>Pinus albicaulis</i>	4	2235	MPB (20%), WPBR (15%)
WBP0924	Siskiyou	Scott River	Marble Mountain	Marble Mountains	<i>Tsuga mertensiana</i>	5	2307	MPB (35%), WPBR (10%)
WBP0925	Siskiyou	n/a	n/a	Big Meadows 1 (Private)	<i>Abies concolor</i>	1	1991	None
WBP0926	Siskiyou	Scott River	Marble Mountain	Big Meadows 2	<i>Tsuga mertensiana</i>	10	2310	Grazing (low), MPB (10%), WPBR (35%)
WBP0927	Siskiyou	Scott River	Marble Mountain	Isinglass Lake 1	<i>Tsuga mertensiana</i>	3	2217	Grazing (medium)
WBP0928	Siskiyou	Scott River	Marble Mountain	Isinglass Lake 2	<i>Pinus albicaulis</i>	8	2182	MPB (10%), WPBR (30%)
WBP0929	Siskiyou	Scott River	Marble Mountain	Above Big Meadows 3	<i>Tsuga mertensiana</i>	6	2309	Grazing (low), MPB (30%)
WBP0930	Siskiyou	Scott River	Marble Mountain	Upper Wright Lake	<i>Pinus albicaulis</i>	12	2379	MPB (10%), WPBR (10%)
WBP0931	Siskiyou	Scott River	Marble Mountain	Red Mountain	<i>Pinus albicaulis</i>	14	2488	None
WBP0932	Siskiyou	Scott River	Marble Mountain	Red Mountain	<i>Pinus albicaulis</i>	6	2366	None

Table 3. Rapid Assessment summary, Klamath National Forest

DbaseID	Site Name	Stand Size	Clumps per hectare	Stems per hectare	Percent Vegetative	Percent Flowering	Percent Fruiting	Mortality by MPB	Mortality by WPBR	Total Mortality	Quality
WBP0900	China Mountain	> 5 acres	75	375	25	0	75	5	0	5	Excellent
WBP0901	China Mountain	> 5 acres	100	700	20	0	80	15	0	15	Excellent
WBP0902	China Mountain	> 5 acres	50	550	50	0	50	40	0	40	Fair
WBP0903	China Mountain	1-5 acres	25	200	50	0	50	0	0	0	Excellent
WBP0904	Goosenest Mountain	> 5 acres	75	425	0	0	100	0	0	0	Excellent
WBP0905	Goosenest Mountain (Private)	> 5 acres	100	625	25	0	75	15	0	15	Excellent
WBP0906	Goosenest Mountain	> 5 acres	100	700	20	0	80	0	0	0	Good
WBP0907	Goosenest Mountain	1-5 acres	50	375	50	0	50	20	0	20	Good
WBP0908	Willow Creek Mountain	1-5 acres	75	425	20	0	80	0	5	5	Excellent
WBP0909	Willow Creek Mountain	1-5 acres	75	625	20	0	80	0	0	0	Excellent
WBP0921	Grizzly Peak	< 1 acre	125	425	40	0	60	0	5	5	Good
WBP0922	Trail Creek	< 1 acre	25	150	50	0	50	0	0	0	Good
WBP0923	Marble Mountains	1-5 acres	75	450	50	0	50	20	0	70% (50% fire, 20% MPB).	Poor
WBP0924	Marble Mountains	1-5 acres	100	600	25	0	75	35	10	45	Fair
WBP0925	Big Meadows 1 (Private)	< 1 acre	0	25	0	0	100	0	0	0	Good
WBP0926	Big Meadows 2	1-5 acres	75	475	25	0	75	10	0	10	Good
WBP0927	Isinglass Lake 1	1-5 acres	0	75	50	0	50	0	0	0	Good
WBP0928	Isinglass Lake 2	1-5 acres	50	375	50	0	50	10	0	10	Good
WBP0929	Big Meadows 3	1-5 acres	100	650	40	0	60	30	0	30	Fair
WBP0930	Upper Wright Lake	1-5 acres	175	1025	30	0	70	10	10	20	Good
WBP0931	Red Mountain	> 5 acres	250	1450	20	0	80	0	0	0	Excellent
WBP0932	Red Mountain	> 5 acres	75	425	25	0	75	0	0	0	Good

# Overview of 2018 Whitebark Pine Surveys, Klamath National Forest




# Russian Wilderness, Salmon River and Scott River Ranger Districts, Klamath NF

**Legend**

**Survey Type, Alliance**


- Rapid Assessment - *Tsuga mertensiana*
- *Pinus albicaulis* Alliance
- *Pinus albicaulis* Presence
- ▭ Ranger District Boundaries
- ▭ Wilderness Boundary



Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community





# Marble Mountain Wilderness, Scott River Ranger Districts, Klamath National Forest




# China Mountain, Scott River Ranger District, Klamath National Forest

**Legend**

**Survey Type, Alliance**

- Rapid Assessment - *Pinus albicaulis*
- Rapid Assessment - *Pinus balfouriana*
- Rapid Assessment - *Abies magnifica*
-  *Pinus albicaulis* Predicted
- Pinus albicaulis* Alliance
- Pinus albicaulis* Presence
- Negative Report
-  National Forest Boundaries




Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community

# Goosenest and Willow Creek Mountain, Goosenest Ranger District, Klamath National Forest

**Legend**

**Survey Type, Alliance**

- Rapid Assessment - Pinus albicaulis
- Rapid Assessment - Abies magnifica
- Rapid Assessment - Tsuga mertensiana
- Pinus albicaulis Predicted
- Pinus albicaulis Alliance
- Pinus albicaulis Presence
- Negative Report
- National Forest Boundaries



Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community