

Whitebark Pine Pilot Fieldwork Report

Modoc National Forest



By Michael Kauffmann¹, Sara Taylor², Kendra Sikes³, and Julie Evens⁴

In collaboration with:

Forrest Guana, Forest Botanist, Modoc National Forest

Diane Ikeda, Regional Botanist, Pacific Southwest Region, USDA Forest Service

January, 2014

November 2018 update: page 46

-
1. Kauffmann, Michael E., Humboldt State University, Redwood Science Project, 1 Harpst Street, Arcata, CA 95521, michaelekauffmann@gmail.com
 2. Taylor, Sara M.
 3. Sikes, Kendra., California Native Plant Society, 2707 K Street, Suite 1, Sacramento, CA 95816, ksikes@cnps.org
 4. Evens, Julie., California Native Plant Society, 2707 K Street, Suite 1, Sacramento, CA 95816, jevans@cnps.org



CALIFORNIA
NATIVE PLANT SOCIETY



Photo on cover page: *Pinus albicaulis* seen from Eagle Peak in the
South Warner Wilderness area, Modoc National Forest

All photos by Michael Kauffmann unless otherwise noted
All figures by Kendra Sikes unless otherwise noted

Acknowledgements: We would like to thank Matt Bokach, Becky Estes, Jonathan Nesmith, Nathan Stephenson, Pete Figura, Cynthia Snyder, Danny Cluck, Marc Meyer, Silvia Haultain, Deems Burton and Peggy Moore for providing field data points or mapped whitebark pine for this project.

Suggested report citation:

Kauffmann, M., S. Taylor, K. Sikes, and J. Evens. 2014. Modoc National Forest: Whitebark Pine Pilot Fieldwork Report. Unpublished report. California Native Plant Society Vegetation Program, Sacramento, CA.

<http://pacslope-conifers.com/conifers/pine/wbp/CNPS-Reports/index.html>

Table of Contents

i	Table of Contents
ii	Figures
ii	Tables
1	Background
3	Introduction
3	Methods and Materials
4	Results
6	Forest-Specific Conclusions, Discussion, and Recommendations
7	Summary Tables from the CNDDDB Rare Plant Occurrence Forms
10	Overview and Regional Detail Maps of 2013 Locations Visited on the National Forest, including negative reports
18	Photos from 2013 Field Work
25	Literature Cited
27	Appendix 1: Inventory and Monitoring Protocols and Field Forms from 2013 including the CNPS Vegetation Rapid Assessment/Relevé Form
40	Appendix 2: Recommended Protocols for Future Work

Figures

- 2 Figure 1: Whitebark pine range in western North America (by Michael Kauffmann)
- 10 Figure 2: Overview Maps of 2013 Locations Visited on the National Forest
- 11 Figure 3: Overview of the Warner Mountain Ranger District
- 12 Figure 4: Whitebark pine in the southern Warner Mountains
- 13 Figure 5: Whitebark pine in the South Warner Wilderness
- 14 Figure 6: Whitebark pine in the central Warner Mountains
- 15 Figure 7: Whitebark pine in the northern Warner Mountains
- 16 Figure 8: Doublehead Ranger District overview
- 17 Figure 9: Doublehead Ranger District whitebark pine negative reports
- 18-19 Figure 10-13: Images of WBP in the southern Warner Mountains
- 19-22 Figures 14-21: Images of WBP in the South Warner Wilderness
- 23 Figures 22-24: Images of WBP in the central Warner Mountains
- 24 Figures 25-27: Images of WBP in the northern Warner Mountains

Tables

- 7 Table 1: Whitebark pine population area by forest in northern California
- 8 Table 2: Rapid assessment summary for Modoc National Forest
- 9 Tables 3: Whitebark pine attributes from Rapid Assessments

Background

Whitebark pine (*Pinus albicaulis*) is a long-lived and slow-growing tree found in upper montane to subalpine forests of southwestern Canada and the western United States. It regularly defines upper treeline and co-occurs with other conifers. Of the approximately 250,000 acres where whitebark pine forms pure stands in California, >95% is on public land, often in remote wilderness settings on National Forest and Park lands; however, the acreage of the pine's presence in mixed-stands across the state is much greater (see Figure 1).

Across the state, the species is found from 1,830 – 4,240 m (6,000'–13,899') in the Sierra Nevada, Cascade, Warner, and Klamath mountains where it is an outlier of a much broader range (Arno et al. 1989, Murray 2005) from the more contiguous Rocky Mountains and Cascades in western North America. Within this range, the species prefers cold, windy, snowy, and generally moist zones. In the moist areas of the Klamath and Cascades, it is most abundant on the warmer and drier sites. In the more arid Warner Mountains and in the Sierra Nevada, the species prefers the cooler north-face slopes and more mesic regions. But some of these phytogeographic patterns are shifting.

Western coniferous forests are currently undergoing large-scale changes in composition and distribution. These changes are due to shifts in the following: climate regimes, insect and fungal pathogen distributions, fire return intervals, fire severity/intensity, and logging practices—among others. High elevation five-needle pines have been harbingers for climate change for millions of years, and because high-elevation ecosystems are likely to be the first to register the impacts of global climate change (Bunn et al. 2005), surveying high elevation five-needle pine is a way to catalog trends in vegetation and climatic shifts.

Unlike other five-needle pines, whitebark pine is set apart in that its seed does not open at maturity and is “wingless”; consequently, they are solely dependent on Clark's nutcrackers (*Nucifraga columbiana*) for seed distribution and future seedling recruitment. The birds open the cone, collect the seeds, and cache them. Inevitably, around 20% of the seeds are forgotten or moved by other animals (Lanner 1996) and, in the years following, clumps of whitebark pine saplings grow from these “forgotten” caches. These two species are keystone mutualist, where the loss of one species would have a profound impact upon the ecosystem as a whole.

Whitebark pine (WBP) is currently the most susceptible of the five-needle pines to mortality due to the combined effects of climate change-induced disturbance. Mortality across much of its range is attributed to white pine blister rust (WPBR) outbreaks caused by the non-native invasive pathogen (*Cronartium ribicola*) (Tomback and Achuff 2010) and native mountain pine beetle (*Dendroctonus ponderosae*) attacks (Logan and Powell 2001, Logan et al. 2010). Decimation to populations in the northern Rocky Mountains has led Canada to list the species as endangered in 2010 (http://www.cosewic.gc.ca/eng/sct1/searchdetail_e.cfm). The current and potential loss of this keystone species in the high mountains of California poses serious threats to biodiversity and losses of ecosystem services, since whitebark pine is one of only a few tree species in these settings.

Mountain pine beetles (MPB) are of concern with respect to high elevation conifers and a warming climate. The beetle is a native insect, having co-evolved with western pine forests in fluctuations of periodic disturbance often followed by cleansing fire regime events. More recently, mass beetle infestations have been correlated with increased climatic warming (Mock 2007). Mountain pine beetles require sufficient thermal input to complete the life cycle in one season. Historically, high elevation ecosystems did not meet these conditions. However, due to recent warming trends, there is adequate thermal input at high elevations for the beetle's lifecycle and infestations of whitebark pine are now increasingly common (Logan and Powell 2001). The preponderance of mass infestations at high elevations has been witnessed throughout California—especially in the arid Warner and eastern Sierra Nevada mountains.

In addition to native insects, a non-native fungal pathogen is affecting high elevation forests. In 1910 white pine blister rust (*Cronartium ribicola*) arrived in a British Columbia port and by 1930 had spread to southern Oregon, infect-

ing western white pine (*Pinus monticola*) and sugar pine (*Pinus lambertiana*) (Murray 2005) along the way. The lifecycle completion requires WPBR to utilize *Ribes* spp. as alternate hosts. In late summer, spores from *Cronartium ribicola* are blown from the *Ribes* host and then enter 5-needle pines through stomata. Upon successful entry, hyphae grow, spread through the phloem, then ultimately swell and kill tissue above the site of infection. Infected trees can survive for over 10 years, but the infection inhibits reproduction (Murray 2005). For species like WBP, which live in fringe habitat and therefore delay reproductive events until conditions are optimal, having an infection that further inhibits cone production is a dangerous proposition. The fungus is found on foxtail and whitebark pines in northwest California (Maloy 2001) where variability in microsite infestation occur (Ettl 2007). On Mount Ashland in the Siskiyou Mountains, blister rust has infected 4 of the 9 WBP trees in the population (Murray 2005). All five-needle native western pines have shown some heritable resistance in the past 100 years (Schoettle et al. 2007), but enduring an infection works against a long-lived pine's survival strategy. Populations of whitebark pine did not evolve to withstand fungal infections.

Seedling establishment for organisms that are on the ecological edge, like WBP, is also jeopardized because of the effects of climate change. Causes of unsuccessful seedling recruitment are many but at high elevation include the effects of fire suppression over the past 100 years. While fire has never been a common phenomenon in high-elevation forests, a shift in fire regime occurred in WBP populations during the Holocene, around 4500 years ago. Before that time fire was not a significant factor in WBP ecology but since has become significant (Murray 2005). The introduction of fire regime suppression in the 1930's is another factor in maintaining whitebark populations. The lack of fire, when coupled with effects of climate change, could also lead to population decline. Whitebark pines need open space for seedling establishment and historically some of this open space has been created by fire events. Fire suppression has also led to increased fire severity and intensity which could be compounded by pathogens. If blister rust and mountain pine beetles continue to move into the high elevations of California, they will potentially generate more dead and downed wood. While considering the potential for the risk of stand replacing fire, this would not mimic historical fire regimes—which have been of low intensity and often focused on individual trees by lightning strikes (Murray 2007).

For more images and discussion of whitebark pine forest health in California see supplementary document (Kauffmann 2014).




Figure 1: Range of WBP in North America (by M. E. Kauffmann)

Introduction

Mapping of whitebark pine occurrence and status/threat has been done primarily using aerial imagery in the National Forests of California by the US Forest Service, including the Pacific Southwest Region - Remote Sensing Lab's CALVEG classification system and maps. The existing USFS vegetation tiles are a result of a 2004-2005 Classification and Assessment with LANDSAT of Visible Ecological Groupings (CALVEG) map product, source imagery ranging from 2002-2009 (USFS 2013c). Even though tile data is continually updated, many stands have not been visited in the field to confirm the accuracy of CALVEG vegetation types. Additionally, little field assessment has been done in the state to identify the presence of whitebark pine, its abundance and status.

The California Native Plant Society (CNPS), working in collaboration with the US Forest Service, initiated field surveys in the summer/fall of 2013 to assess the extent and status of whitebark pine in areas lacking ground surveys in California. Three national forests in the Sierra Nevada and four national forests in the Cascades and Klamath Mountains were selected for field surveys in 2013.

The goals of the field assessments were to verify distribution and status of whitebark pine, ground-truth polygons designated by CALVEG as Whitebark Pine Regional Dominance Type, conduct modified rapid assessments and reconnaissance surveys (recons) on whitebark pine and related stands, and check the USDA Forest Service (USFS) Margins' dataset points for changes in mortality of whitebark pine due to mountain pine beetle and white pine blister rust, if time allowed. Locations within national forests were targeted for the assessment based on potential occurrence of healthy stands in high elevations within the western-most range for the species. Post field assessment, photo interpretation and delineation of whitebark pine extent beyond field surveyed areas were also conducted.

Methods and Materials

The California Native Plant Society (CNPS) obtained existing GIS data from various sources including the USFS Pacific Southwest - Region Remote Sensing Lab's CALVEG maps (USFS 2013c), USFS Forest Health Technology Enterprise Team's National Insect and Disease Risk Model (USFS 2013a) Host species layers, USFS Pacific Southwest Regional Forest Health and Monitoring Aerial Detection Survey Data (USFS 2013b), USFS Forest Health Protection Margins dataset (Bokach 2013), USFS Forest and Inventory Analysis database (USFS 2013d), The Consortium of California Herbaria (UC Berkeley 2013), USFS Central Sierra Province Ecologist-Becky Estes, USFS Southern Sierra Nevada Province Ecologist - Marc Meyer, National Park Service (NPS) Sierra Nevada Network Inventory and Monitoring Program Ecologist - Jonathan Nesmith, US Geological Survey (USGS) Western Ecological Research Center Ecologist - Nathan Stephenson, California Department of Fish and Wildlife (CDFW) Wildlife Biologist - Pete Figura and USFS Northern California Shared Service Center Entomologist - Cynthia Snyder. In addition, we used older sources of whitebark pine distribution in the state for context (Griffin and Critchfield 1972) and for lone populations or individuals not delineated or attributed by CALVEG (Consortium of California Herbaria, 2014).

CNPS also reviewed existing protocols for evaluating whitebark pine vegetation and insect/disease impacts. These protocols included the NPS Standard Operation Procedures for monitoring White Pine (USDOI 2012), Whitebark Pine Ecosystem Foundation (Tomback, et al. 2005), Whitebark Pine Inventory and Monitoring Plot protocol (USFS 2013e) and several government research and staff reports (i.e., Millar et al. 2012, Simons and Cluck 2010, Figura 1997, McKinney et al. 2011, and Maloney et al. 2012). We also discussed the existing protocols for assessing whitebark pine vegetation with USFS staff, including Marc Meyer and Shana Gross.

Upon evaluating existing datasets and obtaining input from local National Forest staff, we identified areas to further ground-truth to better determine the distribution and health/status of whitebark pine on the National Forest lands. Priorities included sampling within wilderness lands and identifying areas with low-levels of insect/disease impact.

We selected the CNPS/CDFW Vegetation Rapid Assessment protocol (see Appendix 2) to gather information

on occurrence, habitat, and impacts of stands with whitebark pine. We modified this protocol to include signs of Mountain Pine Beetle (MPB) and White Pine Blister Rust (WPBR), and overall whitebark pine status/health. The modified rapid assessment aimed to gather as much information on whitebark pine health without spending a significant amount of time establishing plots or collecting data on individual trees. Therefore, the survey technique was stand based to assess the extent of whitebark pine vegetation across broad areas in a short amount of time. Sampling included pure stands, mixed conifer stands, and high elevation krummholz, as long as whitebark pine was deemed a component.

The modifications to the rapid assessment included additional information from Marc Meyer's 'Whitebark Pine Inventory and Monitoring Plot Protocol' such as; whitebark pine impacts from MPB and WPBR, MPB level of attack and % of WBP cones (female). Other protocol information added included; # of individual clumps/stems per area, phenology of WBP (% vegetative, % male flowers and % fruiting) and overall site/ occurrence quality/viability (site + population) from the California Natural Diversity Database (CNDDDB). Since MPB attack and WPBR infestation were the main disturbance of interest to be recorded, USFS Pathologists and Entomologists were contacted for visual aids for proper whitebark pine health assessment. Subsequently, comprehensive field guides were made for recognizing symptoms and signs of MPB and WPBR attack (Kauffmann, 2014).

The reconnaissance (recon) form used for the assessment takes pertinent information from the CNPS/CDFW Vegetation Rapid Assessment protocol to gather simplified, general information about a stand (see Appendix 2). Since the goal of the assessment was to gather information on healthy stands of WBP over a large area, the three purposes of the recon form were to collect data either on 1) WBP stands that were largely diseased or infested, 2) stands attributed as WBP by CALVEG but were incorrect, or 3) WBP stands that were close to stands sampled by a Rapid Assessment.

Areas that were selected for sampling in the Modoc National Forest were based on several approaches including identifying and locating populations that were not yet verified, stand accessibility by road, and wilderness settings that were predicted to have been affected by beetles or rust. These assessment areas were also based on places that Michael Kauffmann had already identified as whitebark pine habitat/population centers (Kauffmann 2013). Danny Cluck and Pete Figura also provided invaluable population details in email communication.

Results

In the Modoc National Forest there are numerous populations of whitebark pine, often isolated on the highest peaks of the Warner Mountains spread in a north-south direction over approximately 60 air miles. The elevational gradient ranges from 2,121-2,998m (7,000-9,892 ft) (which is the summit of Eagle Peak, the highest in the range). For ease of presenting the survey results, the regions reported include: southern Warners, South Warner Wilderness, middle Warners, and north Warners. Within these populations, whitebark pines associate with trees including white fir (*Abies concolor*), lodgepole pine (*Pinus contorta*), western white pine (*Pinus monticola*), ponderosa pine (*Pinus ponderosa* ssp. *ponderosa*), and quaking aspen (*Populus tremuloides*). Though high levels of mortality from mountain pine beetle was observed for several pine species no instances of white pine blister rust were documented.

• Warner Mountain Ranger District

Southern Warner Mountains - Buck Mountain region

This geography of this area is quite similar to that of the Mount Vida-Yellow Mountain region in the northern Warners in that the gentle summits of Buck and Hat Mountains are connected by a high-elevation plateau. The levels and patterns in conifer mortality are similar in both regions as well. Most of the tree mortality was seen in lodgepole (averaging 70%) while whitebark pine mortality in the Buck Mountain region averaged around 20%. The north slopes just below Buck Mountain were being salvage logged for lodgepole pine and grazing of sheep and cattle in this area was common within stands of whitebark pine. There are several mountain summits south of Buck Mountain I recommend for WBP surveys, see recommendation section below. NOTE: Accessibility to this high plateau requires a 4wd vehicle.

South Warner Wilderness

The high-elevation landscape along the spine of the South Warner Wilderness, from approximately Emerson Peak in the south to Squaw Peak in the north, nurtures the largest and most contiguous stands of whitebark pine in northern California. At nearly 8,000 hectares these stands of trees are larger than the WBP acreage found on either the Lassen or the Klamath national forests. Because of the size and varied habitats, the trees are exhibiting a variety of growth and mortality patterns. My surveys into the Eagle Peak/North Emerson Lake region offers only a small glimpse into the lives of whitebark pines in the wilderness.

In the lower elevational limits, north of the Patterson Guard Station as well as on the slopes of summits like Cole and Emerson Peak, the species was exhibiting mortality patterns similar what is seen in the southern and northern Warners. Coupled with high lodgepole pine mortality (common across the Warners) whitebark mortality averaged ~20% on north-facing slopes. However in other areas, downslope expansion through seedling recruitment was common. One area this was observed was along the ridgeline and west-facing meadows between Emerson and Eagle peaks. I hypothesize this is occurring due to fire suppression and large-scale grazing by both sheep (hundreds were seen in this meadow) and cattle. On the north slopes of Eagle Peak, high recruitment was witnessed into north-facing slopes due to decreased snow pack and the new habitat offered to whitebark pine through a longer growing season.

Middle Warners - Bald Mountain region

The area around Cedar and Bald Mountain held the smallest populations, by area, of whitebark pine I surveyed in the Warner Mountains. They were also among the healthiest. Reasons for lack in MPB infestations could include the lower percentage of lodgepole pine which could mitigate the vectoring of mountain pine beetles into the area, the sparse stands of trees which could possibly be less appealing to MPB, the trees often inhabited the xeric ridgelines and west slopes rather than the north slopes (thought this was the preferred habitat on Cedar Mountain), MPB have not yet “found” these trees in large numbers, or the physiological construction of the tree’s resin ducts (Kane et al. 2013). An area that was not surveyed but predicted to hold WBP is to the north of Bald-Cedar Mtns, dissected by FS 118, and includes Buck Mountain (different than southern Warners) and an unnamed plateau just to the south (see map and recommendations for more).

Northern Warners - Mount Vida region

This geography of this area is quite similar to that of the Buck-Hat Mountain region in the southern Warner Mountains in that gentle mountain summits are surrounded and often connected by high elevation, level plateaus. The levels and patterns in conifer mortality are similar as well. While whitebark and lodgepole are common plants in each region, western white pine occur in greater abundance in the northern Warners. Mortality rates for whitebark pine were as high at 70% in certain areas, particularly on the gentle north-facing slopes of Mount Vida but the species composition within the vegetation averaged only ~10% for WBP, with the majority being lodgepole and western white pines, which also had high mortality rates. I recommend surveys around Yellow Mountain and the high plateau branching south and east from Mount Bidwell. I also predict that there is a smaller and more isolated population along the ridge-line south of FS Road 224 in the region around Fandango Peak.

• Doublehead Ranger District

The Medicine Lake highlands were explored including the summit of Mount Hoffman, the highest peak in the region at 2,399m (7,918’ft) but whitebark pine were not found. Forests at the highest elevations included western white pine and mountain hemlock. See negative report map for detailed locations.

Conclusions/Discussion/ Recommendations

The whitebark pine field work in the Modoc National Forest was important in assessing the overall distribution of this vegetation, including significant increases and re-shaping of existing mapped areas of whitebark pine compared to previous delineations from remotely sensing. The increase in mapped area, particularly for the middle Warner Mountains around Bald Mountain was substantial and the size of previously identified population was expanded significantly.

Using the California Natural Diversity Database (CNDDDB) protocol for documenting overall quality and viability of whitebark pine stands observed in the surrounding National Forest areas, we conclude that, overall, populations had fair to excellent viability (probability of persistence) over the next 20 years depending on habitat. The smaller, isolated populations of the middle Warners were excellent but larger more extensive populations, especially on north-facing slopes were often poor to fair in viability. Downslope range expansion through seedling recruitment was common in certain areas of the South Warner Wilderness as well (see results above).

- South Warner Mountains - Buck Mountain region

1. With some of the highest mortality rates seen while ground-truthing the Warner Mountains, it would be advisable to set up several long-term monitoring plots on both Buck and Hat mountains.
2. Ground-truth the extent of WBP's range in this region of Lassen County - especially the area to the south of these highest peaks.
3. Are there whitebark pine on Little Hat Mountain? Elevation suggests they could be there.

- South Warner Wilderness

1. Assimilate all data for the wilderness and create a comprehensive range map for whitebark pine.
2. Monitor the expansion of the species onto north-facing slopes at high elevations and downslope into meadows at the lower extent of the species range within the wilderness.

- Middle Warners - Bald Mountain region

1. Continue to monitor these isolated and small populations of whitebark on both Bald and Cedar mountain. Explore the idea of designating these two peaks and the ridgeline between them as a botanical and geological area.
2. Ground truth the Buck Mountain area (north of Bald Mtn. and different from Buck Mountain previously mentioned) as well as the high ridgeline just south of Buck Mtn. to assess if WBP are present and the extent of the populations.
3. Ground truth Payne Peak, south of 299, to assess if WBP are present and the extent of the populations.

- Northern Warners - Mount Vida region

1. Ground truth the area around Mount Bidwell and Yellow Mountain to assess the health and extent of these whitebark pine populations.
2. Ground truth Fandango Peak and the ridgeline to the north toward Fandango Pass to assess the health and extent of these potential undocumented population centers

Lastly, this report is not comprehensive; it was based upon the available funding, resources and USDA Forest Service staff schedules in 2013. The draft map of whitebark pine distribution (see Figure 2) is therefore not complete but hopefully provides an updated version of whitebark pine distribution from field surveys and aerial interpretation with limited modeled data. The modeled data that is presented from CALVEG in Figure 2 is used to provide areas of data gaps where future field assessments are needed.

More resources for whitebark pine in northern California:

Keeler-Wolf, Todd. 1990. Ecological surveys of FS research natural areas in California.

http://www.fs.fed.us/psw/publications/documents/psw_gtr125/

- o Crater Creek RNA, Klamath
- o Mt. Eddy RNA, Shasta Trinity
- o Sugar Creek RNA, Klamath
- o Antelope Creek Lakes, Klamath

Table 1. Area of whitebark pine populations by national forest region in northern California

Forest	Region	Acres	Hectares
Klamath National Forest	South Goosenest (5 polygons with a small amount in the Shasta-Trinity NF)	2,631	1,065
	North Goosenest (3 polygons)	152	62
	Marble Mountain Wilderness	4,721	1,911
	Russian Wilderness	630	255
	China Mountain Region (some in Shasta-Trinity)	609	246
	total	9,198	3,722
Shasta-Trinity National Forest	Mount Eddy Region	6,048	2,448
	Mount Shasta	11,595	4,692
	Trinity Alps Wilderness	5,671	2,295
	Also see China Mountain and South Goosenest above		
	total	22,039	8,919
Lassen National Forest (including Lassen N.P.)	Within Lassen National Park	11,435	4,628
	Thousand Lakes Wilderness	645	261
	Burney Mountain	15	6
	total	12,095	4,895
Modoc National Forest	Buck Mountain Region	2,401	826
	South Warner Wilderness	20,125	8,548
	Middle Warners	448	181
	North Warners	3,884	1,572
	total	26,858	11,127
Total acreage in the four forest regions of Northern California		70,906	28,633

Table 2. Rapid Assessment summary, Modoc NF

DbaseID	County	Ranger District	Wilderness	Site name	Alliance	Estimated Pct Cover PIAL	Altitude (m)	Impacts
WBP0056	Lassen	Warner Mtn		Southern Warners	Pinus contorta subsp. murrayana	15	2518	Grazing (low), MPB (40%)
WBP0057	Lassen	Warner Mtn		Southern Warners	Pinus albicaulis	2	2548	Grazing (low), MPB (10%)
WBP0058	Lassen	Warner Mtn		Southern Warners	Pinus albicaulis	35	2612	Grazing (low), MPB (10%)
WBP0059	Lassen	Warner Mtn		Southern Warners	Pinus albicaulis	15	n/a	MPB (65%)
WBP0060	Lassen	Warner Mtn		Southern Warners	Populus tremuloides	20	2490	Logging (low), MPB (70%)
WBP0043	Modoc	Warner Mtn		Central Warners	Populus tremuloides	10	2385	Grazing (low)
WBP0044	Modoc	Warner Mtn		Central Warners	Pinus albicaulis	15	2472	Grazing (low), MPB (10%)
WBP0045	Modoc	Warner Mtn		Central Warners	Pinus albicaulis	10	2455	Grazing (low)
WBP0046	Modoc	Warner Mtn		Central Warners	Pinus albicaulis	10	2463	Grazing (low), MPB (5%)
WBP0047	Modoc	Warner Mtn		Central Warners	Pinus monticola	4	2300	Grazing (low)
WBP0048	Modoc	Warner Mtn		Northern Warners	Pinus monticola	8	2367	Grazing (medium)
WBP0049	Modoc	Warner Mtn		Northern Warners	Pinus monticola	4	2400	Grazing (low), MPB (50%)
WBP0050	Modoc	Warner Mtn		Northern Warners	Pinus albicaulis	18	2490	Grazing (low), MPB (5%)
WBP0051	Modoc	Warner Mtn		Northern Warners	Pinus contorta subsp. murrayana	6	2453	MPB (70%)
WBP0052	Modoc	Warner Mtn		Northern Warners	Pinus albicaulis	33	2455	Grazing (low), MPB (5%)
WBP0053	Modoc	Warner Mtn		Northern Warners	Pinus monticola	6	2396	Grazing (low), MPB (10%)
WBP0069	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	18	2370	MPB (40%)
WBP0070	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	23	2424	
WBP0071	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	20	2551	MPB (40%)
WBP0072	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	15	2778	MPB (40%)
WBP0073	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	10	2997	
WBP0074	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	30	2476	MPB (10%)
WBP0075	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	40	2582	MPB (5%)
WBP0076	Modoc	Warner Mtn	South Warner	Warner Wilderness	Pinus albicaulis	10	2366	

Table 3. *Pinus albicaulis* attributes from Rapid Assessments in Modoc NF. N/A is used when data was not recorded.

DbaseID	Site name	Clumps per hectare	Percent Vegetative	Percent Flowering	Percent Fruiting	Mortality by MPB	Total Mortality	Quality
WBP0056	Southern Warners	25.0	25		75	60%	60%	Fair
WBP0057	Southern Warners	30.0	50		50	10%	10%	Good
WBP0058	Southern Warners	30.0	25		75	10%	15%	Good
WBP0059	Southern Warners	30.0	25		75	65%	65%	Poor
WBP0060	Southern Warners	10.0	100			70%	70%	Poor
WBP0043	Central Warners	20.0	40		60	0	0	Excellent
WBP0044	Central Warners	15.0	80		20	10%	10%	Good
WBP0045	Central Warners	n/a	20		80	0	0	Excellent
WBP0046	Central Warners	25.0	40	60		5%	5%	Excellent
WBP0047	Central Warners	5.0	80	20		0	0	Excellent
WBP0048	Northern Warners	10.0	80		20	0	50%	Good
WBP0049	Northern Warners	12.5	80		20	50%	50%	Fair
WBP0050	Northern Warners	50.0	50		50	5%	5%	Good
WBP0051	Northern Warners	15.0	50		50	70%	70%	Poor
WBP0052	Northern Warners	40.0	60		40	5%	5%	Excellent
WBP0053	Northern Warners	25.0	60		40	10%	10%	Good
WBP0069	Warner Wilderness	15.0	100			40%	40%	Fair
WBP0070	Warner Wilderness	20.0	50		50	0	0	Excellent
WBP0071	Warner Wilderness	25.0	50		50	40%	40%	Fair
WBP0072	Warner Wilderness	20.0	60		40	40%	40%	Fair
WBP0073	Warner Wilderness	40.0	80	20		0	0	Excellent
WBP0074	Warner Wilderness	20.0	50		50	10%	10%	Good
WBP0075	Warner Wilderness	30.0	50		50	5%	5%	Excellent
WBP0076	Warner Wilderness	10.0	80		20	0	0	Excellent




Figure 2: Overview Maps of 2013 Locations Visited on the Modoc NF




Figure 3: Warner Mountain Ranger District - Overview




Figure 4: Whitebark Pine in the Warner Mountain Ranger District - southern Warner Mountains




Figure 5: Whitebark Pine in the Warner Mountain Ranger District - South Warner Wilderness




Figure 6: Whitebark Pine in the Warner Mountain Ranger District - central Warner Mountains.




Figure 7: Whitebark Pine in the Warner Mountain Ranger District - northern Warner Mountains.




Figure 8: Doublehead Ranger District - Overview




Figure 9: Doublehead Ranger District - whitebark pine negative reports

Warner Mountain Ranger District - southern Warner Mountains



Figure 10: Looking east toward Hat Mountain from the north flanks of Buck Mountain. Mortality on north slopes averaged ~70% in lodgepole pine (~70%) and ~20% in whitebark pine.



Figure 11: Looking north through relatively healthy whitebark pine on the summit of Buck Mountain.



Figure 12: Mountain pine beetles mortality in lodgepole-whitebark forests between Buck and Hat Mountain

Warner Mountain Ranger District - southern Warner Mountains



Figure 13: Mountain pine beetle infestation sites on the north slopes of Buck Mountain.

Warner Mountain Ranger District - Southern Warner Wilderness



Figure 14: A view of Eagle Peak and high MPB caused mortality in the lodgepole pine forest on its lower east slopes.

Warner Mountain Ranger District - Southern Warner Wilderness



Figure 15: Mountain pine beetles mortality on the north and west slopes of Cole Peak.



Figure 16: Mountain pine beetles mortality on the north and west slopes of Cole Peak - most of the dead trees are whitebark pine.

Warner Mountain Ranger District - Southern Warner Wilderness



Figure 17: Looking southward from the summit of Eagle Peak toward a distant Hat Mountain. The whitebark pine in the front and center of the photograph appear to be recruiting down-slope possibly do to grazing and fire suppression.



Figure 18: Old growth whitebark pine frame the Surprise Valley on the east slopes of Eagle Peak, ~200' below the summit.

Warner Mountain Ranger District - Southern Warner Wilderness



Figure 19: Whitebark pine are recruiting into the south and southeast facing slopes of Eagle Peak due to decreased snowpack and increased growing seasons.



Figure 20: Looking north from Eagle Peak across miles of healthy stands of whitebark pine



Figure 21: New seedling sprouting on the summit of Eagle Peak

Warner Mountain Ranger District - Central Warner Mountains



Figure 22: Looking north to Bald Mountain through a small but healthy stand of WBP



Figure 23: Looking south and east from the summit of Bald Mountain, with a stand of WBP on the east slopes.



Figure 24: Cedar Mountain has an extensive and healthy stand of WBP on its north slopes

Warner Mountain Ranger District - Northern Warner Mountains



Figure 25: Looking east from Mount Vida through MPB induced mortality of whitebark, Lodgepole and western white pine



Figure 26: A saddle just east of Mount Vida with more mixed mortality.



Figure 27: A lone whitebark pine along the Highgrade Trail, looking west toward the Medicine Lake Highlands.

Literature Cited

- Arno, S.F., R.J. Hoff. 1989. Silvics of Whitebark Pine (*Pinus Albicaulis*). USDA Forest Service Technical Report. INT-253.
- BLM. 2014. Federal and State managed lands in California and portions northwest Nevada. Bureau of Land Management, California State Office, Mapping Sciences 5/15/2009. Data available at: <http://www.blm.gov/ca/gis/>
- Bokach, M.. 2013. Margin's dataset. USDA Forest Service, Forest Health Protection Program.
- Bunn, A.G., L.J. Graumlich, D.L Urban. 2005. Trends in twentieth-century tree growth at high elevations in the Sierra Nevada and White Mountains, USA. *The Holocene* 15:481-488.
- CNDDDB. 2014. California Natural Diversity Database (CNDDDB). California Department of Fish and Game, Biogeographic Data Branch, Vegetation Classification and Mapping Program, Sacramento, CA. <http://www.dfg.ca.gov/biogeodata/cnddb/>
- Charlet, D.A. and B. Keimel. 1996 (2012). Atlas of Nevada conifers: a phytogeographic reference. University of Nevada Press, Reno, Nevada. Also, personal communication with author (2012).
- Ettl, G. J. 2007. Ecology of Whitebark Pine in the Pacific Northwest. Proceedings of the Conference Whitebark Pine: A Pacific Coast Perspective USDA Forest Service p20-22
- Figura, P. J. 1997. Structure and dynamics of whitebark pine forests in the South Warner Wilderness, northeastern California. A thesis presented to the faculty of Humboldt State University, Humboldt, CA.
- Greater Yellowstone Whitebark Pine Monitoring Working Group. 2007. Interagency Whitebark Pine Monitoring Protocol for the Greater Yellowstone Ecosystem, Version 1.00. Greater Yellowstone Coordinating Committee, Bozeman, MT <http://www.greateryellowstonescience.org/subproducts/14/72>
- Griffin, J. R., and W.B. Critchfield. 1976. The Distribution of Forest Trees in California. USDA Forest Service: Berkeley, CA.
- Kane, J.M. et al. 2013. Resin duct characteristics associated with tree resistance to bark beetles across lodgepole and limber pines. *Oecologia*. December 2013. <http://dx.doi.org/10.1007/s00442-013-2841-2>
- Kauffmann, M. 2012. Conifer Country: A natural history and hiking guide to the 35 conifers of the Klamath Mountain region. Backcountry Press, Kneeland, California.
- Kauffmann, M. 2013. Conifers of the Pacific Slope. Backcountry Press, Kneeland, California.
- Kauffmann, M. 2014. Whitebark Pine Forest Health in California. Unpublished report. California Native Plant Society Vegetation Program, Sacramento, CA. [<http://pacslope-conifers.com/conifers/pine/wbp/CNPS-Reports/WBP-Forest-Health-California.pdf>]
- Keeler-Wolf, Todd. 1990. Ecological surveys of FS research natural areas in California. Available at: http://www.fs.fed.us/psw/publications/documents/psw_gtr125/
- Lanner, R.M. 1996. Made for each other: A symbiosis of birds and pines. Oxford University Press. New York.
- Little, E.L., Jr. 1971. Atlas of United States trees, volume 1, conifers and important hardwoods: U.S. Department of Agriculture Miscellaneous Publication 1146, 9 p., 200 maps.
- Logan, J.A. and J.A. Powell. 2001. Ghost forests, global warming, and the mountain pine beetle (Coleoptera: Scolytidae). *American Entomologist* (Fall) 47: 160 - 172.
- Logan, J.A., W.W. Macfarlane, and L. Willcox, 2010. Whitebark pine vulnerability to climate-driven mountain pine beetle disturbance in the Greater Yellowstone Ecosystem. *Ecol. Appl.* 20(4): 895–902.
- Maloney, P. E., D.R. Vogler, C.E. Jensen, A.D. Mix. 2012. Ecology of whitebark pine populations in relation to white pine blister rust infection in subalpine forests of the Lake Tahoe Basin, USA: Implications for restoration. *Forest Ecology and Management* 280 (2012) 166–175.
- Maloy, O. C. 2001. White pine blister rust. Online. Plant Health Progress doi:10.1094/PHP-2001-0924-01-HM
- McKinney, S. T., T. Rodhouse, L. Chow, P. Latham, D. Sarr, L. Garrett, L. Mutch. 2011. Long-Term Monitoring of High-Elevation White Pine Communities in Pacific West Region National Parks. USDA Forest Service Proceedings RMRS-P-63.

- Millar, C.I., R. D. Westfall, D. L. Delany, M. J. Bokach. 2012. Forest mortality in high-elevation whitebark pine (*Pinus albicaulis*) forests of eastern California, USA; influence of environmental context, bark beetles, climatic water deficit, and warming. *Canadian Journal of Forest Research* 42: 749–765
- Mock, K.E., et. al. 2007. Landscape-scale genetic variation in a forest outbreak species, the mountain pine beetle (*Dendroctonus ponderosae*). *Molecular Ecology* 16:553-568.
- Murray, M.P. 2005. Our threatened timberlines: the plight of whitebark pine ecosystems. *Kalmiopsis*. 12:25-29
- Murray, M.P. 2007 Fire and Pacific Coast Whitebark Pine. *Proceedings of the Conference Whitebark Pine: A Pacific Coast Perspective* USDA Forest Service p51-60.
- National Park Service (NPS). 2012. Monitoring White Pine (*Pinus albicaulis*, *P. balfouriana*, *P. flexilis*) Community Dynamics in the Pacific West Region - Klamath, Sierra Nevada, and Upper Columbia Basin Networks, Standard Operating Procedures Version 1.0. Natural Resource Report NPS/PWR/NRR—2012/533.
- Sawyer, J.O. 2004. Conifers of the Klamath Mountains. *Vegetation Ecology, Proceedings of the second conference on Klamath-Siskiyou ecology*. 128–135
- Sawyer, J. O. 2006. Northwest California. University of California Press, Berkeley, CA.
- Schoettle, A.W. and R.A. Sniezko. 2007. Proactive intervention to sustain high-elevation pine ecosystems threatened by white pine blister rust. *Journal of Forestry Research*. 12: 327-336.
- Simons, R., D. Cluck. 2010. Whitebark pine monitoring plot protocol for the Warner Mountains, Modoc National Forest. USDA Forest Service, Forest Health Protection and Modoc National Forest.
- Tomback, D.F., and P. Achuff, 2010. Blister rust and western forest biodiversity: ecology, values and outlook for white pines. *Forest Pathology* 40: 186–225.
- Tomback, D. F., R.E. Keane, W. W. McCaughey, C. Smith. 2004. Methods for Surveying and Monitoring Whitebark Pine for Blister Rust Infection and Damage. Whitebark Pine Ecosystem Foundation, Missoula, MT.
- UC Berkeley. 2013. Consortium of California Herbaria. Data provided by the participants of the Consortium of California Herbaria (ucjeps.berkeley.edu/consortium/)
- US. Department of the Interior. 2012. Monitoring White Pine (*Pinus albicaulis*, *P. balfouriana*, *P. flexilis*) Community Dynamics in the Pacific West Region - Klamath, Sierra Nevada, and Upper Columbia Basin Networks, Standard Operating Procedures Version 1.0
- USDA Forest Service. 2013a. Host species layers. U.S. Forest Service Forest Health Technology Enterprise Team; 2013 National Insect and Disease Risk Model. Data available at: <http://www.fs.fed.us/foresthealth/technology/nidrm.shtml>
- USDA Forest Service. 2013b. U.S. Forest Service Pacific Southwest Regional Forest Health Monitoring Aerial Detection Survey Data. Data available at: <http://www.fs.usda.gov/detail/r5/landmanagement/resourcemanagement/?cid=stelprdb5347192>
- USDA Forest Service. 2013c. Vegetation mapping. Existing vegetation (eveg) layers. Pacific Southwest Region Remote Sensing Lab. Data available at <http://www.fs.usda.gov/detail/r5/landmanagement/resourcemanagement/?cid=stelprdb5347192>
- USDA Forest Service. 2013d. Forest and Inventory Analysis database. Forest Inventory and Analysis National Program. Data available at: <http://www.fia.fs.fed.us/tools-data/>
- USDA Forest Service. 2013e. Whitebark Pine Inventory and Monitoring Plot Protocol. USFS Region 5 Ecology Program and Forest Health Protection Program.

Appendix 1: Inventory and Monitoring Protocols and Field Forms from 2013

CALIFORNIA NATIVE PLANT SOCIETY / DEPARTMENT OF FISH AND GAME PROTOCOL FOR COMBINED VEGETATION RAPID ASSESSMENT AND RELEVÉ SAMPLING FIELD FORM

(Modified for WBP)

July 8, 2013

Introduction

This protocol describes the methodology for both the relevé and rapid assessment vegetation sampling techniques as recorded in the combined relevé and rapid assessment field survey form dated June 28, 2013. The same environmental data are collected for both techniques. However, the relevé sample is plot-based, with each species in the plot and its cover being recorded. The rapid assessment sample is based not on a plot but on the entire stand, with 12-20 of the dominant or characteristic species and their cover values recorded. For more background on the relevé and rapid assessment sampling methods, see the relevé and rapid assessment protocols at www.cnps.org.

Selecting stands to sample:

To start either the relevé or rapid assessment method, a stand of vegetation needs to be defined.

A stand is the basic physical unit of vegetation in a landscape. It has no set size. Some vegetation stands are very small, such as alpine meadow or tundra types, and some may be several square kilometers in size, such as desert or forest types.

A stand is defined by two main unifying characteristics:

- 1) It has compositional integrity. Throughout the site, the combination of species is similar. The stand is differentiated from adjacent stands by a discernable boundary that may be abrupt or indistinct.
- 2) It has structural integrity. It has a similar history or environmental setting that affords relatively similar horizontal and vertical spacing of plant species. For example, a hillside forest originally dominated by the same species that burned on the upper part of the slopes, but not the lower, would be divided into two stands. Likewise, sparse woodland occupying a slope with very shallow rocky soils would be considered a different stand from an adjacent slope with deeper, moister soil and a denser woodland or forest of the same species.

The structural and compositional features of a stand are often combined into a term called homogeneity. For an area of vegetated ground to meet the requirements of a stand, it must be homogeneous (uniform in structure and composition throughout).

Stands to be sampled may be selected by evaluation prior to a site visit (e.g., delineated from aerial photos or satellite images), or they may be selected on site during reconnaissance (to determine extent and boundaries, location of other similar stands, etc.).

Depending on the project goals, you may want to select just one or a few representative stands of each homogeneous vegetation type for sampling (e.g., for developing a classification for a vegetation mapping project), or you may want to sample all of them (e.g., to define a rare vegetation type and/or compare site quality between the few remaining stands).

For the rapid assessment method, you will collect data based on the entire stand.

Selecting a plot to sample within in a stand (for relevés only):

Because many stands are large, it may be difficult to summarize the species composition, cover, and structure of an entire stand. We are also usually trying to capture the most information as efficiently as possible. Thus, we are typically forced to select a representative portion to sample.

When sampling a vegetation stand, the main point to remember is to select a sample that, in as many ways possible, is representative of that stand. This means that you are not randomly selecting a plot; on the contrary, you are actively using your own best judgment to find a representative example of the stand.

Selecting a plot requires that you see enough of the stand you are sampling to feel comfortable in choosing a representative plot location. Take a brief walk through the stand and look for variations in species composition and in stand structure. In many cases in hilly or mountainous terrain look for a vantage point from which you can get a representative view of the whole stand. Variations in vegetation that are repeated throughout the stand should be included in your plot. Once you assess the variation within the stand, attempt to find an area that captures the stand's common species composition and structural condition to sample.

Plot Size

All relevés of the same type of vegetation to be analyzed in a study need to be the same size. Plot shape and size are somewhat dependent on the type of vegetation under study. Therefore, general guidelines for plot sizes of tree-, shrub-, and herbaceous communities have been established. Sufficient work has been done in temperate vegetation to be confident the following conventions will capture species richness:

Herbaceous communities: 100 sq. m plot

Special herbaceous communities, such as vernal pools, fens: 10 sq m plot

Shrublands and Riparian forest/woodlands: 400 sq. m plot

Open desert and other shrublands with widely dispersed but regularly occurring woody species: 1000 sq. m plot

Upland Forest and woodland communities: 1000 sq. m plot

Plot Shape

A relevé has no fixed shape, though plot shape should reflect the character of the stand. If the stand is about the same size as a relevé, the plot boundaries may be similar to that of the entire stand. If we are sampling streamside riparian or other linear communities, our plot dimensions should not go beyond the community's natural ecological boundaries. Thus, a relatively long, narrow plot capturing the vegetation within the stand, but not outside it would be appropriate. Species present along the edges of the plot that are clearly part of the adjacent stand should be excluded.

If we are sampling broad homogeneous stands, we would most likely choose a shape such as a circle (which has the advantage of the edges being equidistant to the center point) or a square (which can be quickly laid out using perpendicular tapes).

Definitions of fields in the protocol

Relevé or Rapid Assessment: Circle the method that you are using.

I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION

Polygon/Stand #: Number assigned either in the field or in the office prior to sampling. It is usually denoted with a four-letter abbreviation of the sampling location and then a four-number sequential number of that locale (e.g. CARR0001 for Carrizo sample #1). The maximum number of letters/numbers is eight.

Air photo #: The number given to the aerial photo in a vegetation-mapping project, for which photo interpreters have already done photo interpretation and delineations of polygons. If the sample site has not been photo-interpreted, leave blank.

Date: Date of the sampling.

Name(s) of surveyors: The full names of each person assisting should be provided for the first field form for the day. On successive forms, initials of each person assisting can be recorded. Please note: The person recording the data on the form should circle their name/initials.

GPS waypoint #: The waypoint number assigned by a Global Positioning System (GPS) unit when marking and storing a waypoint for the sample location. Stored points should be downloaded in the office to serve as a check on the written points and to enter into a GIS.

For relevé plots, take the waypoint in the southwest corner of the plot or in the center of a circular plot.

GPS name: The name/number assigned to each GPS unit. This can be the serial number if another number is not assigned.

Datum: (NAD 83) The standard GPS datum used is NAD 83. If you are using a different datum, note it here.

Bearing, left axis at SW pt (note in degrees) of Long or Short side: For square or rectangular plots: from the SW corner (= the GPS point location), looking towards the plot, record the bearing of the axis to your left. If the plot is a rectangle, indicate whether the left side of the plot is the long or short side of the rectangle by circling "long" or "short" side (no need to circle anything for circular or square plots). If there are no stand constraints, you would choose a circular or square plot and straight-sided plots should be set up with boundaries running in the cardinal directions. If you choose a rectangular plot that is not constrained by the stand dimensions, the short side should run from east to west, while the long side should run from north to south.

UTM coordinates: Easting (UTME) and northing (UTMN) location coordinates using the Universal Transverse Mercator (UTM) grid. Record in writing the information from a GPS unit or a USGS topographic map.

UTM zone: Universal Transverse Mercator zone. Zone 10 is for California west of the 120th longitude, zone 11 is for California east of 120th longitude, which is the same as the straight portion of California's eastern boundary.

Error ±: The accuracy of the GPS location, when taking the UTM field reading. Please record the error units by circling feet (ft), meters (m), or positional dilution of precision (pdop). If your GPS does not determine error, insert N/A in this field.

Is GPS within stand? Yes / No Circle “Yes” to denote that the GPS waypoint was taken directly within or at the edge of the stand being assessed for a rapid assessment, or circle “No” if the waypoint was taken at a distance from the stand (such as with a binocular view of the stand).

If No, cite from waypoint to stand, distance (note in meters) & bearing (note in degrees): An estimate of the number of meters and the compass bearing from the GPS waypoint to the stand.

Elevation: Recorded from the GPS unit or USGS topographic map. Please circle feet (ft) or meters (m).

Photograph #s: Write the name or initials of the camera owner, JPG/frame number, and direction of photos (note the roll number if using film). Take four photos in the main cardinal directions (N, E, S, W) clockwise from the north, from the GPS location. If additional photos are taken in other directions, please note this information on the form. Also include overview photos of Whitebark pine.

Stand Size: Estimate the size of the entire stand in which the sample is taken. As a measure, one acre is about 4000 square meters (approximately 64 x 64 m), or 208 feet by 208 feet. One acre is similar in size to a football field.

Plot Size: If this is a relevé, circle the size of the plot.

Plot Shape: Record the length and width of the plot and circle measurement units (i.e., ft or m). If it is a circular plot, enter radius (or just put a check mark in the space).

Exposure: (Enter actual ° and circle general category): With your back to the general uphill direction of the slope (i.e., by facing downhill of the slope), read degrees of the compass for the aspect or the direction you are standing, using degrees from north, adjusted for declination. Average the reading over the entire stand, even if you are sampling a relevé plot, since your plot is representative of the stand. If estimating the exposure, write “N/A” for the actual degrees, and circle the general category chosen. “Variable” may be selected if the same, homogenous stand of vegetation occurs across a varied range of slope exposures. Select “all” if stand is on top of a knoll that slopes in all directions or if the same, homogenous stand of vegetation occurs across all ranges of slope.

Steepness: (Enter actual ° and circle general category): Read degree slope from a compass or clinometer. If estimating, write “N/A” for the actual degrees, and circle the general category chosen.. Make sure to average the reading across the entire stand even if you are sampling in a relevé plot.

Topography: First assess the broad (Macro) topographic feature or general position of the stand in the surrounding watershed, that is, the stand is at the top, upper (1/3 of slope), middle (1/3 of slope), lower (1/3 of slope), or bottom. Circle all of the positions that apply for macrotopography.

Then assess the local (Micro) topographic features or the lay of the area (e.g., surface is flat or concave). Circle only one of the microtopographic descriptors.

Geology: Geological parent material of site. If exact type is unknown, use a more general category (e.g., igneous, metamorphic, sedimentary). See code list for types.

Soil Texture: Record soil texture that is characteristic of the site (e.g., coarse loamy sand, sandy clay loam). See soil texture key and code list for types.

Upland or Wetland/Riparian (circle one): Indicate if the stand is in an upland or a wetland. There are only two options. Wetland and riparian are one category. Note that a site need not be officially delineated as a wetland to qualify as such in this context (e.g., seasonally wet meadow).

% Surface cover (abiotic substrates). It is helpful to imagine “mowing off” all of the live vegetation at the base of the plants and removing it – you will be estimating what is left covering the surface. The total should sum to 100%. Note that non-vascular cover (lichens, mosses, cryptobiotic crusts) is not estimated in this section.

% Water: Estimate the percent surface cover of running or standing water, ignoring the substrate below the water.

% BA Stems: Percent surface cover of the plant basal area, i.e., the basal area of stems at the ground surface. Note that for most vegetation types BA is 1-3% cover. Estimate for a set area (e.g., 400 m²) of BA to help calibrate on this % (on average % is between 1.5-4.5% for conifers)

% Litter: Percent surface cover of litter, duff, or wood on the ground.

% Bedrock: Percent surface cover of bedrock.

% Boulders: Percent surface cover of rocks > 60 cm in diameter.

% Stone: Percent surface cover of rocks 25-60 cm in diameter.

% Cobble: Percent surface cover of rocks 7.5 to 25 cm in diameter.

% Gravel: Percent surface cover of rocks 2 mm to 7.5 cm in diameter.

% Fines: Percent surface cover of bare ground and fine sediment (e.g. dirt) < 2 mm in diameter.

% Current year bioturbation: Estimate the percent of the sample or stand exhibiting soil disturbance by fossorial organisms (any organism that lives underground). Do not include disturbance by ungulates. Note that this is a separate estimation from surface cover.

Past bioturbation present? Circle Yes if there is evidence of bioturbation from previous years.

% Hoof punch: Note the percent of the sample or stand surface that has been punched down by hooves (cattle or native grazers) in wet soil.

Fire Evidence: Circle Yes if there is visible evidence of fire, and note the type of evidence in the “Site history, stand age and comments section,” for example, “charred dead stems of *Quercus berberidifolia* extending 2 feet above resprouting shrubs.” If you are certain of the year of the fire, put this in the Site history section.

Site history, stand age, and comments: Briefly describe the stand age/seral stage, disturbance history, nature and extent of land use, and other site environmental and vegetation factors. Examples of disturbance history: fire, landslides, avalanching, drought, flood, animal burrowing, or pest outbreak. Also, try to estimate year or frequency of disturbance. Examples of land use: grazing, timber harvest, or mining. Examples of other site factors: exposed rocks, soil with fine-textured sediments, high litter/duff build-up, multi-storied vegetation structure, or other stand dynamics.

Disturbance code / Intensity (L,M,H): List codes for potential or existing impacts on the stability of the plant community. Characterize each impact each as L (=Light), M (=Moderate), or H (=Heavy). For invasive exotics, divide the total exotic cover (e.g. 25% *Bromus diandrus* + 8% *Bromus madritensis* + 5% *Centaurea melitensis* = 38% total exotics) by the total % cover of all the layers when added up (e.g. 15% tree + 5% low tree + 25% shrub + 40% herbs = 85% total) and multiply by 100 to get the % relative cover of exotics (e.g. 38% total exotics/85% total cover = 45% relative exotic cover). L = 0-33% relative cover of exotics; M = 34-66% relative cover, and H = > 66% relative cover. See code list for impacts.

List percent of WBP impacted by Mountain Pine Beetle (39-MPB/L/approx. % impacted) and White Pine Blister Rust (40-WPBR/H/approx. % impacted) within the stand. For Mountain Pine Beetle, search the bole for entry holes (reddish colored pitch) or frass. For WPBR, search for ‘signs’ of an active canker (i.e., a canker with visible aecia, or fruiting bodies containing spores), or ‘symptoms’ of any of the following five indicators: rodent chewing, flagging, swelling, roughened bark, and oozing sap. Explain signs and symptoms in the notes and take photos when necessary.

II. HABITAT AND VEGETATION DESCRIPTION

California Wildlife-Habitat Relationships (CWHR)

For CWHR, identify the size/height class of the stand using the following tree, shrub, and/or herbaceous categories. These categories are based on functional life forms.

Tree DBH: Circle one of the tree size classes provided when the tree canopy closure exceeds 10 percent of the total cover, or if young tree density indicates imminent tree dominance. Size class is based on the average diameter at breast height (dbh) of each trunk (standard breast height is 4.5ft or 137cm). When marking the main size class, make sure to estimate the mean diameter of all trees over the entire stand, and weight the mean if there are some larger tree dbh's. The "T6 multi-layered" dbh size class contains a multi-layered tree canopy (with a size class T3 and/or T4 layer growing under a T5 layer and a distinct height separation between the classes) exceeding 60% total cover. Stands in the T6 class need also to contain at least 10% cover of size class 5 (>24" dbh) trees growing over a distinct layer with at least 10% combined cover of trees in size classes 3 or 4 (>11-24" dbh). This is weighted: In your representative area add number of trees for each category and record above (T1,T2,T3, etc). Can square root later to get the weighted average for this category (if there are many sizes).

Shrub: Circle one of the shrub size classes provided when shrub canopy closure exceeds 10 percent (except in desert types) by recording which class is predominant in the survey. Shrub size class is based on the average amount of crown decadence (dead standing vegetation on live shrubs when looking across the crowns of the shrubs).

Herb: Circle one of the herb height classes when herbaceous cover exceeds 2 percent by recording the predominant class in the survey. Note: This height class is based on the average plant height at maturity, not necessarily at the time of observation.

Desert Palm/Joshua Tree: Circle one of the palm or Joshua tree size classes by averaging all the stem-base diameters (i.e. mean diameter of all stem-base sizes). Diameter is measured at the plant's base above the bulge near the ground.

Desert Riparian Tree/Shrub: Circle one of the size classes by measuring mean stem height (whether tree and/or shrub stand).

Overall Cover of Vegetation

Provide an estimate of cover for the following categories below (based on functional life forms). Record a specific number for the total aerial cover or "bird's-eye view" looking from above for each category, estimating cover for the living plants only. Litter/duff should not be included in these estimates. The porosity of the vegetation should be taken into consideration when estimating percent cover (how much of the sky can you see when you are standing under the canopy of a tree, or how much light passes through the canopy of the shrub layer?).

To come up with a specific number estimate for percent cover, first use generalized cover classes as reference aids such as the CWHR cover classes (<2%, 2-9%, 10-24%, 25-39%, 40-59%, 60-100%) or the modified Braun-Blanquet cover-abundance scale (<1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, >75%). While keeping these intervals in mind, you can then refine your estimate to a specific percentage for each category below.

% Total NonVasc cover: The total cover of all lichens, bryophytes (mosses, liverworts, hornworts), and cryptogammic crust on substrate surfaces including downed logs, rocks and soil, but not on standing or inclined trees or vertical rock surfaces.

% Total Vasc Veg cover: The total cover of all vascular vegetation taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute vegetation cover, disregarding overlap of the various tree, shrub, and/or herbaceous layers and species. Could use densitometer to calibrate, but sometimes this provides an over-estimate.

% Cover by Layer

% Conifer Tree /Hardwood Tree: The total foliar cover (considering porosity) of all live tree species, disregarding overlap of individual trees. Estimate conifer and hardwood covers separately.

Please note: These cover values should not include the coverage of regenerating tree species (i.e., tree seedlings and saplings).

% Regenerating Tree: The total foliar cover of seedlings and saplings, disregarding overlap of individual recruits. See seedling and sapling definitions below.

%Shrub: The total foliar cover (considering porosity) of all live shrub species disregarding overlap of individual shrubs.

%Herbaceous: The total cover (considering porosity) of all herbaceous species, disregarding overlap of individual herbs.

Height Class by Layer

Modal height for conifer tree /hardwood tree, shrub, and herbaceous categories: Provide an estimate of height for each category listed. Record an average height value per each category by estimating the mean height for each group. Please use the following height intervals to record a height class: 01 = < 1/2m, 02=1/2-1m, 03 = 1-2 m, 04 = 2-5 m, 05 = 5-10 m, 06 = 10-15 m, 07 = 15-20 m, 08 = 20-35 m, 09 = 35-50 m, 10 => 50m.

Species List and Coverage

- If mistletoe present add in what species it is living on
- Record absolute percent cover of dead tree species (can include saplings and seedlings)

For rapid assessments, list the 10-20 species that are dominant or that are characteristically consistent throughout the stand. These species may or may not be abundant, but they should be constant representatives in the survey. When different layers of vegetation occur in the stand, make sure to list species from each stratum. As a general guide, make sure to list at least 1-2 of the most abundant species per stratum.

For relevés, list all species present in the plot, using the second species list page if necessary.

For both sample types, provide the stratum:

T = Tree. A woody perennial plant that has a single trunk.

S = Shrub. A perennial, woody plant, that is multi-branched and doesn't die back to the ground every year.

H = Herb. An annual or perennial that dies down to ground level every year.

E = SEedling. A tree species clearly of a very young age that is < 1" dbh.

A = SApling. 1" - <6" dbh and young in age, OR small trees that are < 1" diameter at breast height, are clearly of appreciable age, and kept short by repeated browsing, burning, or other disturbance.

N = Non-vascular. Includes moss, lichen, liverworts, hornworts, cryptogamic crust, and algae.

Be consistent and don't break up a single species into two separate strata. The only time it would be appropriate to do so is when one or more tree species are regenerating, in which case the Seedling and/or Sapling strata should be recorded for that species. These may be noted on the same line, e.g.:

Strata	Species	%Cover	C
T/E/A	Quercus douglasii	40/<1/<1	

If a species collection is made, it should be indicated in the collection column with a "C" (for collected). If the species is later keyed out, cross out the species name or description and write the keyed species name in pen on the data sheet. Do not erase what was written in the field, because this information can be used if specimens get mixed up later. If the specimen is then thrown out, the "C" in the collection column should be crossed out. If the specimen is kept but is still not confidently identified, add a "U" to the "C" in the collection column (CU = collected and unconfirmed). In this case the unconfirmed species epithet should be put in parentheses [e.g. *Hordeum (murinum)*]. If the specimen is kept and is confidently identified, add a "C" to the existing "C" in the collection column (CC = Collected and confirmed).

Use Jepson Manual nomenclature. Write out the genus and species of the plant. Do not abbreviate. When uncertain of an identification (which you intend to confirm later) use parentheses to indicate what part of the determination needs to be confirmed. For example, you could write out *Brassica (nigra)* if you are sure it is a Brassica but you need further clarification on the specific epithet.

Provide the % absolute aerial cover for each species listed. When estimating, it is often helpful to think of coverage in terms of the following cover intervals at first:

<1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, >75%.

Keeping these classes in mind, then refine your estimate to a specific percentage. All species percent covers may total over 100% because of overlap.

Include the percent cover of snags (standing dead) of trees and shrubs. Note their species, if known, in the "Stand history, stand age and comments" section.

For rapid assessments, make sure that the major non-native species occurring in the stand also are listed in the space provided in the species list with their strata and % cover. For relevés, all non-native species should be included in the species list.

Also for relevés, you can record the <1% cover in two categories: r = trace (i.e., rare in plot, or solitary individuals) and + = <1% (few individuals at < 1% cover, but common in the plot).

Unusual species: List species that are locally or regionally rare, endangered, or atypical (e.g., range extension or range limit) within the stand. This field will be useful to the Program for obtaining data on regionally or locally significant populations of plants.

INTERPRETATION OF STAND

Field-assessed vegetation alliance name: Name of alliance or habitat following the most recent CNPS classification system or the Manual of California Vegetation (Sawyer J.O., Keeler-Wolf T., and Evens, J. 2009). Please use scientific nomenclature, e.g., *Quercus agrifolia* forest. An alliance is based on the dominant or diagnostic species of the stand, and is usually of the uppermost and/or dominant height stratum. A dominant species covers the greatest area. A diagnostic species is consistently found in some vegetation types but not others.

Please note: The field-assessed alliance name may not exist in the present classification, in which case you can provide a new alliance name in this field. If this is the case, also make sure to state that it is not in the MCV under the explanation for “Confidence in alliance identification.”

Field-assessed association name (optional): Name of the species in the alliance and additional dominant/diagnostic species from any strata, as according to CNPS classification. In following naming conventions, species in differing strata are separated with a slash, and species in the uppermost stratum are listed first (e.g., *Quercus douglasii*/*Toxicodendron diversilobum*). Species in the same stratum are separated with a dash (e.g., *Quercus lobata*-*Quercus douglasii*).

Please note: The field-assessed association name may not exist in the present classification, in which you can provide a new association name in this field.

Adjacent Alliances/direction: Identify other vegetation types that are directly adjacent to the stand being assessed by noting the dominant species (or known type). Also note the distance away in meters from the GPS waypoint and the direction in degrees aspect that the adjacent alliance is found

(e.g., *Amsinckia tessellata* / 50m, 360°N *Eriogonum fasciculatum* /100m, 110°).

Confidence in Identification: (L, M, H) With respect to the “field-assessed alliance name”, note whether you have L (=Low), M (=Moderate), or H (=High) confidence in the interpretation of this alliance name.

Explain: Please elaborate if your “Confidence in Identification” is low or moderate. Low confidence can occur from such things as a poor view of the stand, an unusual mix of species that does not meet the criteria of any described alliance, or a low confidence in your ability to identify species that are significant members of the stand.

Phenology: Indicate early (E), peak (P) or late (L) phenology for each of the strata.

Other identification problems or mapping issues: Discuss any further problems with the identification of the assessment or issues that may be of interest to mappers. Note if this sample represents a type that is likely too small to map. If it does, how much of the likely mapping unit would be comprised of this type. For example: “this sample represents the top of kangaroo rat precincts in this general area, which are surrounded by vegetation represented by CARR000x; this type makes up 10% of the mapping unit.” Depending on who mapped polygon (Calveg, etc); we should denote that information here.

Is polygon >1 type: Yes / No (circle one): In areas that have been delineated as polygons on aerial photographs/imagery for a vegetation-mapping project, assess if the polygon is mapped as a single stand. “Yes” is noted when the polygon delineated contains the field-assessed alliance and other vegetation type(s), as based on species composition and structure. “No” is noted when the polygon is primarily representative of the field-assessed alliance.

If yes, explain: If “Yes” above, explain the other vegetation alliances that are included within the polygon, and explain the amount and location that they cover in the polygon.

Other CNDDDB/Whitebark Pine (WBP) monitoring Data:

Trees/stems are assessed within a representative portion of the stand (using a specific radius or area for averaging).

Mountain Pine Beetle (MPB) Level: Should equal 100%.

Note the level of mountain pine beetle attack using the following:

- 0 = No evidence of attack or beetle pitch tubes or unknown
- 1 = less than 5 observable beetle pitch tubes ('hits')
- 2 = less than 50% of the bole is attacked; sporadic pitch tubes spread on most parts of the bole or several localized areas with a high density (>10) pitch tubes
- 3 = greater than 50% of the bole is attacked; numerous pitch tubes spread on many parts of the bole

% of WBP Cones (female only): Should equal 100%.

Record the number of cones in the tree/stem using the following numeric system:

- 0 = no cones
- 1 = 1 to 10 cones
- 2 = 11 to 100 cones
- 3 = greater than 100 cones

Total # WBP individuals or clumps and size (CNDDDB):

The number of individuals observed/detected during assessment. This should be recorded as clumps (or # of stems within # of clumps) per defined area (square meters, hectares, acres, etc.).

Phenology of WBP (CNDDDB): Should equal 100%.

The average percent of WBP that is vegetative, flowering (nascent female cones) and/or fruiting (mature female cones).

% WBP mortality:

These percentages are for mortality of trees/stems from mountain pine beetle (MPB) or white pine blister rust (WPBR); 'Other' can be % mortality from both MPB and WPBR; including WPBR mortality on other species E.g. WPBR-PIMO/PIBA 5% (white pine blister rust on *Pinus monticola* or *Pinus balfouriana* at 5% cover) or unknown causes.

Overall site/occurrence quality/viability (site + population) (CNDDDB):

Is the likely persistence of the occurrence into the future Excellent, Good, Fair, or Poor? This is an assessment of the overall viability of this occurrence. Both the quality & condition of the site and of the occurrence must be considered when scoring. Take into account population size, demography, viability over time, site condition, and any disturbances. And also see additional characteristics at: <http://www.natureserve.org/explorer/eorankguide.htm>

Determination of WBP: Please indicate how the species identification was determined.

CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)

Relevé or Rapid Assessment (circle one) (Revised June 28, 2013)

For Office Use:	Final database #:	Final vegetation type name:	Alliance Association
-----------------	-------------------	-----------------------------	----------------------

I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION

Polygon/Stand #:	Air photo:	Date:	Name(s) of surveyors (circle recorder):
------------------	------------	-------	---

GPS wypt #: _____ GPS name: _____ Datum: _____ or NAD83. Bearing, left axis at SW pt _____ (degrees) of Long / Short side
 UTMN _____ UTMN _____ Zone: 10 / 11 (circle one) Error: ± _____ ft / m / pdop
 GPS within stand? Yes / No If No, cite from waypoint to stand, distance _____ (meters) & bearing _____ (degrees)

Elevation: _____ ft / m Camera Name/Photograph #'s: _____

Stand Size (ac/ha): <1, 1-5, >5 ac | _____ ha Plot Size (m²): 10 / 100 / 400 | Plot Shape _____ x _____ m or Circle Radius _____ m
 Exposure, Actual °: _____ NE NW SE SW Flat Variable All | Steepness, Actual °: _____ 0° 1-5° 5-25° > 25

Topography: Macro: top upper mid lower bottom | Micro: convex flat concave undulating
 Geology code: _____ Soil Texture code: _____ | Upland or Wetland/Riparian (circle one)

% Surface cover: (Incl. outcrops) (>60cm diam) (25-60cm) (7.5-25cm) (2mm-7.5cm) (Incl sand, mud)
 H20: _____ BA Stems: _____ Litter: _____ Bedrock: _____ Boulder: _____ Stone: _____ Cobble: _____ Gravel: _____ Fines: _____ =100%

% Current year bioturbation _____ Past bioturbation present? Yes / No | Fire evidence: Yes / No (if yes, explain below)
 Habitat description, surrounding land use, comments (CNDDDB):

Disturbance / Intensity (L,M,H) _____ / _____ / _____ / _____ WBP Impact 39 / _____ / _____ 40 / _____ / _____

II. HABITAT AND VEGETATION DESCRIPTION

Tree DBH: **T1** (<1" dbh), **T2** (1-6" dbh), **T3** (6-11" dbh), **T4** (11-24" dbh), **T5** (>24" dbh), **T6** multi-layered (T3 or T4 layer under T5, >60% cover)
 Shrub: **S1** seedling (<3 yr. old), **S2** young (<1% dead), **S3** mature (1-25% dead), **S4** decadent (>25% dead)
 Herbaceous: **H1** (<12" plant ht.), **H2** (>12" ht.) % NonVasc cover: _____ % Vasc Veg cover: _____
 % Cover - Conifer tree / Hardwood tree: _____ / _____ Regenerating Tree: _____ Shrub: _____ Herbaceous: _____
 Height Class - Conifer tree / Hardwood tree: _____ / _____ Regenerating Tree: _____ Shrub: _____ Herbaceous: _____
 Height classes: 01=<1/2m 02=1/2-1m 03=1-2m 04=2-5m 05=5-10m 06=10-15m 07=15-20m 08=20-35m 09=35-50m 10=>50m

Species, Stratum, and % cover. Stratum categories: T=Tree, S = Shrub, H= Herb, E = SEedling, A = SApling, N= Non-vascular.
 % cover intervals for reference: <1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, 75%.

Strata	Species	% dead	% cover	C	Strata	Species	% dead	% cover	C

Other rare taxa in stand (CNDDDB) _____

III. INTERPRETATION OF STAND

Field-assessed vegetation alliance name: _____

Field-assessed association name (optional): _____

Adjacent alliances/direction: _____ / _____, _____ / _____

Confidence in alliance identification: L M H Explain: _____

Other identification or mapping information: _____ Phenology (E,P,L): Herb _____ Shrub _____ Tree _____

Is poly >1 type: Yes / No If yes, explain: _____

**CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)
Other CNDDDB/Whitebark Pine Monitoring Data:**

Polygon/Stand #:
MBP Level: 0=no attack _____ % 1=>5 hits _____ % 2=<50% of bole attacked _____ % 3=>50% of bole attacked _____ %
Avg % of WBP Cones: No cones _____ % 1-10 cones _____ % 11-100 _____ % >100 _____ %
Total # individuals or clumps (WBP) and size (CNDDDB) _____ # per _____ hectares
Phenology of WBP (CNDDDB): Vegetative _____ % Flowering (cones) _____ % Fruiting (cones) _____ %
%WBP mortality: MPB _____ % WPBR _____ % Other: _____ % _____ %
Overall site/occurrence quality/viability (site + population) (CNDDDB): <input type="checkbox"/> Excellent <input type="checkbox"/> Good <input type="checkbox"/> Fair <input type="checkbox"/> Poor
Determination of WBP: Keyed _____ By another person (name) _____ Compared with photo/drawing _____ Other _____

Sample Rapid Assessment from the Boulder Peak Region - Marble Mountain Wilderness

**CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)
Other CNDDDB/Whitebark Pine Monitoring Data:**

Polygon/Stand #: BP01
MBP Level: 0=no attack <u>70</u> % 1=>5 hits <u>—</u> % 2=<50% of bole attacked <u>—</u> % 3=>50% of bole attacked <u>30</u> %
Avg % of WBP Cones: No cones <u>70</u> % 1-10 cones <u>20</u> % 11-100 <u>10</u> % >100 <u>—</u> %
Total # individual clumps (WBP) and size (CNDDDB) <u>15</u> # per <u>2</u> hectares
Phenology of WBP (CNDDDB): vegetative <u>70</u> % flowering (cones) <u>20</u> % fruiting (cones) <u>20</u> %
%WBP mortality: MPB <u>30</u> % WPB <u>0</u> % Other: <u>—</u> % _____ %
Overall site/occurrence quality/viability (site + population) (CNDDDB): <input type="checkbox"/> Excellent <input type="checkbox"/> Good <input checked="" type="checkbox"/> Fair <input type="checkbox"/> Poor
Determination of WBP: Keyed _____ By another person (name) _____ Compared with photo/drawing _____ Other _____

White fir / Shasta fir ATTACK by fir engraver with 20% mortality

WWP (P. monticola) 1 dead MPB.

CNPS and CDFG Combined Vegetation Rapid Assessment and Relevé Field Form (modified for WBP project)
 Relevé or Rapid Assessment (circle one) (Revised June 28, 2013)

For Office Use: Final database #: Final vegetation type name: Alliance Association

I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION

Polygon/Stand #: BPO1 Air photo: Date: 9/20/13 Name(s) of surveyors (circle recorder): KARKMAN

GPS wypt #: GPS name: Datum: or NAD83. Bearing, left axis at SW pt (degrees) of Long / Short side
 UTME 492814 UTMN 4604449 Zone: 10 / 11 (circle one) Error: ±2.3 ft / m / pdop
 GPS within stand? Yes / No If No, cite from waypoint to stand, distance (meters) & bearing (degrees)

Elevation: 1985 ft / m Camera Name/Photograph #'s:

Stand Size (ac/ha): <1, 1-5, >5 ac ha Plot Size (mz): 10 / 100 / 500 Plot Shape x m or Circle Radius 12 m
 Exposure, Actual °: NE NW SE SW Flat Variable All | Steepness, Actual °: 3° 0° 1-5° 5-25° >25

Topography: Macro: top upper mid lower bottom | Micro: convex flat concave undulating
 Geology code: GABDO Soil Texture code: Upland or Wetland/Riparian (circle one)

% Surface cover: (Incl. overcrops) (>60cm diam) (25-60cm) (7.5-25cm) (2mm-7.5cm) (Incl sand, mud)
 H20: 50 BA Stems: 30 Litter: 57 Bedrock: Boulder: 10 Stone: 5 Cobble: 5 Gravel: 5 Fines: 10 =100%

% Current year bioturbation Past bioturbation present? Yes / No | Fire evidence: Yes No (if yes, explain below)
 Habitat description, surrounding land use, comments (CNDDDB):

On edge of creek, 1/2 mile below lower Wright Lake (on trail)
 meadow edge - RIPARIAN. Fire evidence from 20+ years ago
 now with white fir encroaching.

Disturbance / Intensity (L,M,H) Grazing / L WBP Impact 39 1M, 50% 40 16, 30%

II. HABITAT AND VEGETATION DESCRIPTION

Tree DBH: T1 (<1" dbh), T2 (1-6" dbh), T3 (6-11" dbh), T4 (11-24" dbh), T5 (>24" dbh), T6 multi-layered (T3 or T4 layer under T5, >60% cover)
 Shrub: S1 seedling (<3 yr. old), S2 young (<1% dead), S3 mature (1-25% dead), S4 decadent (>25% dead)
 Herbaceous: H1 (<12" plant ht.), H2 (>12" ht.) % NonVasc cover: 35 % Vasc Veg cover: 65
 % Cover - Conifer tree / Hardwood tree: 15 / Regenerating Tree: 5 Shrub: 25 Herbaceous: 20
 Height Class - Conifer tree / Hardwood tree: 05 / Regenerating Tree: 02 Shrub: 02 Herbaceous: 01
 Height classes: 01=<1/2m 02=1/2-1m 03=1-2m 04=2-5m 05=5-10m 06=10-15m 07=15-20m 08=20-35m 09=35-50m 10=>50m

Species, Stratum, and % cover. Stratum categories: T=Tree, S=Shrub, H=Herb, E=SEedling, A=SApling, N=Non-vascular.
 % cover intervals for reference: <1%, 1-5%, >5-15%, >15-25%, >25-50%, >50-75%, 75%

Strata	Species	% dead	% cover	C	Strata	Species	% dead	% cover	C
T	Whitebark	30%	5%		S	Servicberry		3%	
T	White fir	10	10%		S	Ribes sp.		4%	
T	Shasta fir		3%		H	Grass sp.		10%	
T	Western white pine (i)	dead-1 year			H	Worm sp		5%	
S	SALIX sp.		5%		H	elderberry		5%	
S	Snowberry		5%		T	FOXTAIL pine - 4 new		3%	
S	VACINIUM sp.		5%			reg trees			
S	Sorbus sp.		3%						

Other rare taxa in stand (CNDDDB)

III. INTERPRETATION OF STAND

Field-assessed vegetation alliance name: White fir - WBP
 Field-assessed association name (optional): meadow to
 Adjacent alliances/direction: EAST
 Confidence in alliance identification: L (M) H Explain:

Other identification or mapping information: Along trail Phenology (E,P,L): Herb Shrub Tree

Is poly >1 type: Yes / No If yes, explain:

Appendix 2: Recommended Protocols for Future Work

Whitebark Pine Inventory and Monitoring Plot Protocol

Revised: May 16, 2013

Introduction:

This protocol was developed collaboratively by the USFS Region 5 Ecology Program and Forest Health Protection Program to provide inventory and status-and-trend monitoring data in stands dominated by whitebark pine (*Pinus albicaulis*) or lodgepole pine (*P. contorta*) with whitebark pine as a codominant species. It also focuses on stands that have experienced recent tree mortality related to insects (mountain pine beetle) or diseases (white pine blister rust). This protocol was developed using elements of the Regional Ecology Program post-fire regeneration monitoring protocol and the Forest Health Protection Whitebark Pine Monitoring Plot Protocol for the Warner Mountains, Modoc National Forest (the FHP protocol is based on *Greater Yellowstone Whitebark Pine Monitoring Working Group [GYWPMWG] Interagency Whitebark Pine Monitoring Protocol for the Greater Yellowstone Ecosystem 2007*).

I. Site attributes:

- Record the location (geographic or watershed scale), site (topographic scale), and plot number (micro-scale).
- Use a GPS device to locate plot center - take care to avoid biasing the location.
- Monument permanent plots (established for monitoring rather than inventory) with 2-foot long rebar driven approximately 1.5 ft into the ground at plot center. Label with plot number and mount safety cap. Include brief notes of plot location using distinctive landscape features, if any.
- Record the date that data were collected.
- Record the crew names of the people collecting the data.
- Establish a plot with a radius of 12.6 m, which is approximately 0.05 ha (0.124 acres). Flag four places around the perimeter for reference.
- Record the dominant tree species present.
- Take one photograph from a point 12.6 m south of the plot center, looking north. Make sure you have something (pin flag) at plot center so it can be relocated. using the photo. Take another photograph from a point 12.6 m north of the plot center, looking south (toward pin flag). Record both photo numbers.
- Record the average slope of the plot in percent (use clinometer).
- Record the average aspect of the plot in degrees (use compass [make sure you have the right declination])

II. Vegetation and ground cover attributes

- Estimate the cover (%) of: *basal vegetation* (i.e. the area covered by the bases of tree boles, shrub stems, herbs), *litter*, *bare ground*, *rock* (>2 mm diameter), and *woody debris* (>3 inches [7.5 cm] diameter), summing to 100% (imagine chopping off all vegetation at ground level, what do you have left?; it is rare for basal vegetation to be more than 5%, unless there are trees or many large shrubs in the plot). Record cover values to nearest 5%, using 0.5% as trace cover
- Estimate vegetation cover to nearest 1% (1-10% cover), 5% (10-30%) or 10% (30-100%):

- % Overstory veg cover = cover of plants >2 m in height (trees and tall shrubs; this is a snapshot of total canopy cover taken from above, i.e. it is the % of the plot that has trees/tall shrubs covering it. Tree/tall shrubs growing completely beneath other trees/tall shrubs are not counted as they cannot be seen from above) (see Figure 1)
 - Estimate cover of live trees and tall shrubs
 - Also estimate % dead cover (trace circles around the remnants of dead trees >2 m in height). When this value is added to the live cover it should give us an estimate of the total pre-beetle mortality live cover.
- % Shrub and Herbaceous plant cover = cover of understory vegetation <2 m in height (this is a snapshot of total understory cover taken from above, i.e. it is the % of the plot that has understory vegetation covering it. Understory plants growing completely beneath other plants are not counted, as they cannot be seen from above) (see Figure 1).
- Record separately the cover of aspen <3 m in height. Aspen >3 m height should be recorded as overstory cover.




Fig. 1. Circular plot, with four species of understory plants (colored). The blue crosshairs are added to aid in estimating cover. The understory vegetation cover is about 64% (the total plot area minus the area that is not covered by live vegetation). The gray species (shrub) has 49% cover, the dark blue species (shrub) has 17% cover, the light blue species (grass) has 4%, and the orange species (forb) has 6% (each orange circle is 1% in this case). Due to plant overlap, summing the different species' cover values gives a value that is larger than the total understory vegetation cover (76% vs. 64%). Overall shrub cover in this plot is 58% (two shrub species, subtracting overlap; summed up [i.e., ignoring overlap], the two species have 66% cover between them). Herb cover is 9.5%; forb cover is 6%; grass cover is 4%. Each shrub species will have its own cover entered in the species-cover section of the datasheet. Cover is measured by drawing a line around the outside of the plant canopy, ignoring gaps that may be found within the perimeter. For plots of this size (500 m²), your clipboard is about 0.015% of the plot area.

III. Basal area, snags, and litter depth

- Use the basal area gauge (20 factor) to record the basal area of live and dead whitebark pine and other tree species (e.g., lodgepole pine) in the stand
 - Swinging the gauge around the plot center, tally the number of trees that are larger than the 20 factor aperture. Count live and dead trees separately for each species.
- Record the species and dbh of any older snags (>7 years; **prior to 2005**) in the plot. Only record snags that are >1.37 m tall.
- Measure litter depth at 3 locations midpoint between plot center and plot perimeter in 3 directions (0°, 120°, and 240°).

IV. Tree regeneration attributes

- Tally the number of seedlings and saplings (trees less than 7.6 cm dbh of each tree species for each age class)
 - Use a separate row for each species and basal cluster (see below).
 - Count the number of live and dead stems arising from each seedling or sapling cluster. Clusters are defined using the following two criteria:
 - Stems are less than 10 cm at the base from the cluster of other stems
 - Diameter of stem (saplings only) must not exceed 25% of next largest stem in the cluster
 - Determine minimum age by counting the bud scars, subtracting the current year
 - Record dbh for saplings (>1.37 m height) only
 - In the Health Code column, note the number of seedlings or saplings in each cluster that exhibit health issues and include the appropriate health codes for these numbers (e.g., 2-a). Health codes include:
 - C = cankers or stem swelling
 - SC = stalactiform canker (*P. contorta* only) as spindle-like in middle of bole
 - P = pitching
 - F = branch flagging
 - S = needle spots
 - T = twig beetle sign (e.g. terminal branch flagging and pitch tubes)
 - 2 = secondary beetle
 - M = dwarf mistletoe
 - R = native rusts – take photo and collect sample when available
 - H = sapsucker/woodpecker holes
 - A = aecia (i.e., rust fruiting body) or aeciospores
 - Take closeup photos of any branches displaying aecia and consider collecting samples for laboratory identification
 - Aecia could be a sign on WPBR or a native rust
 - Record the height for the tallest individual seedling of each species

V. Understory vegetation attributes

- Measure the modal height and overall cover for the four most common (by % cover) shrub and herbaceous plant species in the plot. Additional understory species may be noted in the Notes section. Especially note the presence of species in the genera *Ribes*, *Castilleja*, and *Pedicularis* (WPBR secondary hosts)
 - Measure cover to nearest 5%, 0.5% = trace cover
 - Modal height is the most common height

VI. Notes section

Items of interest to record in the notes section:

- If fire scars or other evidence of fire are in the plot
- If plot is located on a unique (non-granitic) substrate (e.g., pumice soils)
- If plot has been treated in some way - specify
- If non-native species are on plot or adjacent to plot - specify
- If other mortality agents (insects, diseases) are present - specify
- If WPBR, mountain pine beetle, twig beetle, or other potential mortality agents are observed adjacent to plot but not recorded within the plot
- If conifer stumps are present from trees that may have parented seedlings before they were cut
- Additional understory species if more than four shrubs or herbaceous plants
- Other notes?

VII. Tree attributes

For all trees (>7.6 cm dbh) record the following information

- Species ID, and number live and dead stems in each cluster. Tree clusters are defined by stems that are less than 1 m at the base from the cluster of other stems of similar size (diameter and height).
- Individual stems growing in close proximity will be defined as individual tree stems or branches using the following GYWPMWG (2007) criteria:
 - 1) There must be a discernible growth groove that separates that stem from other stems of the tree.
 - 2) The diameter of a given stem must be more than 25% of the diameter of the largest stem.
 - 3) The stem must be less than one foot from the “mother” tree to which it is associated. Otherwise it is to be considered as a separate seedling, sapling, or tree.
 - 4) The angle of the stem in question must be no less than a 45° angle from the main stem.
- For each cluster, provide a consecutively-numbered cluster ID number. For each stem within a cluster, provide a stem ID value.
- Record the dbh of all live and dead stems in each cluster.

- For monumented monitoring plots, nail aluminum tags to all live trees that are counted with the basal area gauge (20 factor) sweep. Begin consecutively-numbered tags at the northernmost tree proceeding clockwise. Nail tags at dbh so that each tag faces plot center, leaving approximately one inch of space between the nail head and the tree bole in order to avoid damage during tree growth. Enter tag numbers in the Notes column.
- Note with a checkmark whether live basal sprouts are present for a given tree cluster. These basal sprouts are defined as smaller-diameter (typically <7.6 cm dbh) live stems located at the base and often surrounding clusters of larger live and dead tree clusters (typically with stems exceeding 20 cm dbh).
- Note the % of live crown in the stem (largest live is the default) in increments of 10% using the following coding system: 1 = 10%, 2 = 20%, 3 = 30%, etc.
- Note the level of mountain pine beetle attack using the following:
 - 0 = No evidence of attack or beetle pitch tubes or unknown
 - 1 = less than 5 observable beetle pitch tubes ('hits')
 - 2 = less than 50% of the bole is attacked; sporadic pitch tubes spread on most parts of the bole or several localized areas with a high density (>10) pitch tubes
 - 3 = greater than 50% of the bole is attacked; numerous pitch tubes spread on many parts of the bole
- Estimate the time since mountain pine beetle attack based on the following system:
 - 0 = less than one year since attack (occurred during current season); typically little sign of crown discoloration or dead needles but evidence of beetle attack
 - 1 = approximately one year since attack (last season); crown shows significant density of dead or dying needles (substantial portion of crown contains brown/orange colored needles)
 - 2 = two years since attack; entire crown consists of dead (brown/orange colored) needles that are mostly intact
 - 3 = three years since attack; most but not all of dead needles have fallen from crown, with few clusters of dead needles retained
 - 4 = four to seven years since attack; no dead needles retained in crown; smaller branches may have broken off and fallen, with most larger to medium branches retained
- Record the number of cones in the tree using the following numeric system:
 - 0 = no cones
 - 1 = 1 to 10 cones
 - 2 = 11 to 100 cones
 - 3 = greater than 100 cones
- Record the health code for each tree using the codes listed under the tree regeneration attributes section (see above)
- In Notes column record any remarkable observations pertaining to a tree or tree cluster, including:
 - lightning or fire scars

- evidence of other damage caused by wildlife, humans, or unknown causes
- Other notable features or observations

VIII. Seed-Caching Wildlife Point Counts (optional, if time permits)

- At end of vegetation sampling period, return to each plot and spend 5 minutes noting any visual or auditory sign of Clark's Nutcracker (*Nucifraga columbiana*), Douglas' squirrel (*Tamiasciurus douglasii*), lodgepole chipmunk (*Neotamias speciosus*), golden-mantled ground squirrel (*Callospermophilus lateralis*), or any other seed-eating species within 50 m of each side of transect. Record plot number, observer, time and date of survey, and number of each species observed at each sample point (i.e., plot).
- Note any observations of seed caching, seed dispersal, or seed predation during survey period.
- Record data on separate field notebook

2018 Update

Whitebark Pine Pilot Fieldwork Report

Modoc National Forest



By Michael Kauffmann¹, Jennifer J. Buck-Diaz, Raphaela Floreani Buzbee,
Julie Evens, and Sydney Magner

In collaboration with:
Heidi Guenther, Forest Botanist, Modoc National Forest
Diane Ikeda, Regional Botanist, Pacific Southwest Region, USDA Forest Service
NOVEMBER 2018

1. Kauffmann, Michael E., PO Box 777, Bayside, CA 95524, michaelekauffmann@gmail.com



CALIFORNIA
NATIVE PLANT SOCIETY



ABOVE: Bark beetle mortality on Mount Bidwell.

Methods and Materials

The California Native Plant Society (CNPS) obtained updated GIS data from various sources including the USFS Pacific Southwest - Region Remote Sensing Lab's CALVEG maps (USFS 2018), and California Department of Fish and Wildlife (CDFW) - High Elevation Species Sightings Sierra Nevada (CDFW 2018). In addition, we used the updated species profile from the CNPS Rare Plant Program, which maps the distribution of whitebark pine using many sources such as herbarium specimens and other observational data (CNPS 2018).

CNPS reviewed the CNPS/CDFW Vegetation Rapid Assessment protocol used in the 2013 pilot project for evaluating whitebark pine vegetation and insect/disease impacts. We made minor edits and clarification to the data sheet and protocol in order to collect standardized data across the five forests. A reconnaissance (recon) form was used to gather simplified, general information about stands of vegetation, especially for WBP stands that were near or adjacent to stands already sampled by a full Rapid Assessment.

Notes on Presence and Alliance

A vegetation alliance is a category of vegetation classification which describes repeating patterns of plants across a landscape. Each alliance is defined by plant species composition, and reflects the effects of local climate, soil, water, disturbance, and other environmental factors. Alliances are commonly used in vegetation mapping.

Presence indicates that whitebark pine can be found but is not the dominant vegetation pattern.



An even-aged cohort regenerating in a stand impacted by mountain pine beetles.

The following areas, prioritized for future work after our 2013 surveys, were visited in the summer of 2018.

Northern Warners - Mount Vida region



1. Ground truth the area around Mount Bidwell and Yellow Mountain to assess the health and extent of these whitebark pine populations.
2. Ground truth Fandango Peak and the ridgeline to the north toward Fandango Pass to assess the health and extent of these potential undocumented population centers.

The range of whitebark pine was much less extensive than predicted in our 2013 mapping and early surveys. Most of the survey time was spent around Yellow Mountain and Mount Bidwell as well as other small, isolated summits above 7,500 feet. While whitebark did occur on all these summits, the predicted presence was not as extensive—though whitebark alliance was found in almost all locations.

Mount Vida was visited in 2013 and found to host a large, extensive stand of whitebark pine. Based on this, predictions for extensive stands were made for north of here. The only area where this was found to be the case in 2018 was along the high elevation ridgeline of Mount Bidwell. Along this 3 mile ridgeline, whitebark pine alliance was strong on the north slopes and presence extended downslope as lodgepole pine became the alliance. The 2006-2012 beetle attacks were evident with mortality in older whitebark pines approaching 50% in some areas, but there was no further evidence of mountain pine beetles in the past ~5 years. Sapling regeneration was also seen and often vigorous.

Between Mount Bidwell and Mount Vida, whitebark pine were found scattered on the highest summits. The most extensive of these stands was on the west slopes of Yellow Mountain. Across this large area, mountain pine beetles killed up to 70% of the lodgepole pine but whitebark pine mortality was much less, 20% on average.

Along the unnamed ridgeline south of Fandango Pass, only a handful of individual whitebark pines were found. All were generally healthy with some signs of mortality in lodgepole pine from the 2006-2012 beetle attack.



Middle Warners - Bald Mountain region

1. Ground truth the Buck Mountain area (north of Bald Mtn.) as well as the high ridgeline just south of Buck Mtn. to assess if WBP are present and the extent of the populations.

On this survey day CNPS was joined by the Modoc National Forest Botanist Heidi Guenther. We began by surveying Buck Mountain in the central Warner Mountains. No whitebark pines were found. The ridge-line was flat and extensive, exceeding 7,500 feet across 10+ acres, but whitebark pine were absent.

From this area, we moved south to survey the northern part of Bald Mountain where we found several small stands of whitebark pine alliance not located in 2013. In these stands, trees showed >5% mortality.





Table 1: Area by forest as mapped by Michael Kauffmann.

Forest		2014 Presences (acres)	2018 Presences (acres)	2018 Alliance (acres)
Klamath National Forest	total	9,198	5,708.6	659.5
Shasta-Trinity National Forest	total	22,039	Did not map for alliance in 2013, only presence. Not revised in 2018.	
Lassen National Forest (including Lassen N.P.)	total	12,095		
Modoc National Forest	total	26,858	23,472.6	1,357.5
Total acreage in the four forests of Northern California		70,906	63,315	--

Notes on the area changes:

Because of the general average high elevation in the northern Warner Mountains, near the Oregon border, I predicted the presence of whitebark pine to be more extensive after our 2013 surveys. I only surveyed Mount Vida that year and the trees were common in the region around the summit. This did not translate to the north. In fact, most of the populations of whitebark pine north of Mount Vida were small stands (5-10 acres) of WBP alliance. The exception was along the extensive ridgeline of Mount Bidwell, where a vast stand of trees occurs.



Small stands of Pinus albicaulis alliance persist on several unnamed summits north of Mount Vida.



Yellow Mountain - Pinus albicaulis is restricted to the west slopes where it forms an alliance.



Mount Bidwell -- the most extensive stands of Pinus albicaulis in the northern Warner Mountains.

Recommendations for future work:

- South Warner Mountains - Buck Mountain region

1. With some of the highest mortality rates seen while ground-truthing the Warner Mountains, it would be advisable to set up several long-term monitoring plots on both Buck and Hat mountains.
2. Are there whitebark pine on Little Hat Mountain? Elevation suggests they could be there.

- South Warner Wilderness

1. Assimilate all data for the wilderness and create a comprehensive range map for whitebark pine.
2. Monitor the expansion of the species onto north-facing slopes at high elevations and downslope into meadows at the lower extent of the species range within the wilderness.

- Middle Warners - Bald Mountain region

1. Continue to monitor these isolated and small populations of whitebark on both Bald and Cedar mountain. Explore the idea of designating these two peaks and the ridgeline between them as a botanical and geological area.
2. Ground truth Payne Peak, south of 299, to assess if WBP are present and the extent of the populations.



Small stands on Bald Mountain as seen in the picture below.



Modoc National Forest staff joined CNPS for surveys on Bald Mountain.


Table 2. Rapid Assessment summary, Modoc National Forest

DbaseID	County	Ranger District	Wilderness	Site Name	Alliance	Estimated Pct Cover PIAL	Altitude (m)	Impacts
WBP0910	Modoc	Warner Mountain	n/a	Yellow Mountain	<i>Pinus monticola</i>	3	2376	None
WBP0911	Modoc	Warner Mountain	n/a	Yellow Mountain	<i>Pinus albicaulis</i>	13	2433	MPB (5%)
WBP0912	Modoc	Warner Mountain	n/a	Yellow Mountain	<i>Pinus contorta ssp. murrayana</i>	3	2420	MPB (15%)
WBP0913	Modoc	Warner Mountain	n/a	Yellow Mountain	<i>Pinus albicaulis</i>	10	2457	None
WBP0914	Modoc	Warner Mountain	n/a	Mt. Bidwell	<i>Pinus albicaulis</i>	11	2513	Grazing (low), MPB (75%)
WBP0915	Modoc	Warner Mountain	n/a	Mt. Bidwell	<i>Pinus contorta ssp. murrayana</i>	7	2476	None
WBP0917	Modoc	Warner Mountain	n/a	North Warners	<i>Artemisia tridentata</i>	3	2343	None
WBP0918	Modoc	Warner Mountain	n/a	North Warners	<i>Pinus monticola</i>	7	2349	None
WBP0919	Modoc	Warner Mountain	n/a	Bald Mountain	<i>Pinus monticola</i>	4	2366	None
WBP0920	Modoc	Warner Mountain	n/a	Bald Mountain	<i>Pinus albicaulis</i>	10	2467	None

Table 3. Rapid Assessment summary, Modoc National Forest

DbaseID	Site name	Stand size	Clumps per hectare	Stems per hectare	Percent Vegetative	Percent Flowering	Percent Fruiting	Mortality by MPB	Mortality by WPBR	Total Mortality	Quality
WBP0910	Yellow Mountain	1-5 acres	0	75	50	0	50	0	0	0	Excellent
WBP0911	Yellow Mountain	1-5 acres	200	1200	20	0	80	5	0	5	Excellent
WBP0912	Yellow Mountain	1-5 acres	25	225	20	0	80	15	0	15	Fair
WBP0913	Yellow Mountain	> 5 acres	50	375	20	0	80	0	0	0	Excellent
WBP0914	Mt. Bidwell	> 5 acres	50	375	20	0	80	75	0	75	Fair
WBP0915	Mt. Bidwell	1-5 acres	75	525	10	0	90	0	0	0	Fair
WBP0917	Northern Warners	< 1 acre	0	25	0	0	100	0	0	0	Excellent
WBP0918	Northern Warners	1-5 acres	25	150	20	0	80	0	0	0	Excellent
WBP0919	Bald Mountain	< 1 acre	100	500	25	0	75	0	0	0	Excellent
WBP0920	Bald Mountain	1-5 acres	200	1125	10	0	90	0	0	0	Excellent

Overview of Whitebark Pine Surveys, Warner Mountain Ranger District, Modoc NF










2018 Whitebark Pine Surveys, Warner Mountain Ranger District, Modoc NF

Legend

2018 Surveys

Survey Type, Alliance

-  Rapid Assessment - *Pinus albicaulis*
-  Rapid Assessment - *Pinus contorta* ssp. *murrayana*
-  Rapid Assessment - *Pinus monticola*
-  Rapid Assessment - *Artemisia tridentata*
-  *Pinus albicaulis* Predicted
-  National Forest Boundaries




Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community

Mount Bidwell, Yellow Mountain and Mount Vida area, Warner Mountain Ranger District, Modoc NF

Legend

Survey Type, Alliance

- Rapid Assessment - *Pinus albicaulis*
- Rapid Assessment - *Pinus contorta* ssp. *murrayana*
- Rapid Assessment - *Pinus monticola*
- Rapid Assessment - *Artemisia tridentata*
- Pinus albicaulis* Predicted
- Pinus albicaulis* Alliance
- Pinus albicaulis* Presence
- Negative Report
- National Forest Boundaries









Sources: Esri, HERE, Garmin, Intermap, increment P Corp., GEBCO, USGS, FAO, NPS, NRCAN, GeoBase, IGN, Kadaster NL, Ordnance Survey, Esri Japan, METI, Esri China (Hong Kong), swisstopo, © OpenStreetMap contributors, and the GIS User Community

Bald and Cedar Mountains, Warner Mountain Ranger District, Modoc NF

Legend

Survey Type, Alliance

-  Rapid Assessment - *Pinus albicaulis*
-  Rapid Assessment - *Pinus monticola*
-  *Pinus albicaulis* Predicted
-  *Pinus albicaulis* Alliance
-  *Pinus albicaulis* Presence
-  Negative Report

